

For Research Use Only

# Dystroglycan Monoclonal ANTIBODY



Catalog Number: 66735-1-Ig **1 Publications**

## Basic Information

<b>Catalog Number:</b> 66735-1-Ig	<b>GenBank Accession Number:</b> BC012740	<b>Purification Method:</b> Protein G purification
<b>Size:</b> 150UL, Concentration: 1000 µg/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 1605	<b>CloneNo.:</b> 2B1G12
<b>Source:</b> Mouse	<b>Full Name:</b> dystroglycan 1 (dystrophin-associated glycoprotein 1)	<b>Recommended Dilutions:</b> WB 1:5000-1:50000 IHC 1:500-1:2000
<b>Isotype:</b> IgG1	<b>Calculated MW:</b> 97 kDa	
<b>Immunogen Catalog Number:</b> AG27222	<b>Observed MW:</b> 43 kDa, 30 kDa	

## Applications

### Tested Applications:

FC, IHC, WB, ELISA

### Cited Applications:

FC

### Species Specificity:

Human, Mouse, Rat, Pig

### Cited Species:

human

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB:** A549 cells, mouse brain tissue, pig brain tissue, rat brain tissue, NCI-H1299 cells, HeLa cells, HepG2 cells

**IHC:** mouse heart tissue, mouse skeletal muscle tissue

## Background Information

Dystroglycan, also known as DAG1 or DG, was originally isolated from skeletal muscle as an integral membrane component of the dystrophin-glycoprotein complex (DGC). In addition to skeletal muscle, dystroglycan is strongly expressed in heart and smooth muscle, as well as many non-muscle tissues including brain and peripheral nerve (PMID: 12556455). The dystroglycan is involved in a number of processes including laminin and basement membrane assembly, sarcolemmal stability, cell survival, peripheral nerve myelination, nodal structure, cell migration, and epithelial polarization. Dystroglycan consists of two subunits (alpha and beta), which are translated from a single mRNA as a propeptide that is proteolytically cleaved into two noncovalently associated proteins (PMID: 16410545). Alpha-dystroglycan is a 156-kDa extracellular peripheral glycoprotein, while beta-dystroglycan is a 43-kDa transmembrane protein (PMID: 9858474). The 43-kDa beta-dystroglycan can be cleaved into a ~30-kDa form (PMID: 14678802; 18458097; 17255331).

## Notable Publications

Author	Pubmed ID	Journal	Application
Nicolás Sarute	32719120	Proc Natl Acad Sci U S A	FC

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com  
W: ptglab.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

