For Research Use Only

## AQP1 Monoclonal antibody

Catalog Number:66805-1-lg 4 Publications





Basic Information	Catalog Number: 66805-1-lg	GenBank Accession Number: BC022486	Purification Method: Protein G purification	
	Size:	GenelD (NCBI):	CloneNo.:	
	150ul, Concentration: 1400 ug/ml b		2E5D6	
	Nanodrop and 1000 ug/ml by Bradformethod using BSA as the standard;	<sup>ord</sup> UNIPROT ID: P29972	Recommended Dilutions: WB 1:5000-1:20000	
	Source:	Full Name: aquaporin 1 (Colton blood group)	IHC 1:200-1:800	
	Mouse		IF-P 1:200-1:800	
	lsotype: lgG1	Calculated MW: 269 aa, 29 kDa		
	Immunogen Catalog Number: AG14093	Observed MW: 28 kDa, 38-40 kDa		
Applications	Tested Applications:	Positive C	Positive Controls: WB : human heart tissue, human skeletal muscle tissue, pig kidney tissue	
	WB, IHC, IF-P, ELISA Cited Applications:			
	IF		an kidney tissue,	
	Species Specificity: Human, pig		an kidney tissue,	
	Cited Species: human, mouse			
	Note-IHC: suggested antigen TE buffer pH 9.0; (*) Alternat retrieval may be performed v	ively, antigen		
	buffer pH 6.0			
Background Information	AQP1 is a member of aquaporins (A AQP1 is expressed in most tissues i	n the mammalian body. Alterations r. The predicted molecular weight o	•	
	AQP1 is a member of aquaporins (A AQP1 is expressed in most tissues i variety of diseases, including cance glycosylated form can also be obser	n the mammalian body. Alterations r. The predicted molecular weight o	ning proteins facilitating water transpor of AQP1 expression have been linked to AQP1 is around 28 kDa, while highly AQP1cation	
	AQP1 is a member of aquaporins (A AQP1 is expressed in most tissues i variety of diseases, including cance glycosylated form can also be obser Author Pu	n the mammalian body. Alterations r. The predicted molecular weight or ved around 40-45 kDa. (1530176)	of AQP1 expression have been linked to AQP1 is around 28 kDa, while highly	
	AQP1 is a member of aquaporins (A AQP1 is expressed in most tissues i variety of diseases, including cance glycosylated form can also be obser Author Pu Lili Huang 39	n the mammalian body. Alterations r. The predicted molecular weight of ved around 40-45 kDa. (1530176) bmed ID Journal	of AQP1 expression have been linked to AQP1 is around 28 kDa, while highly AQP1cation	
	AQP1 is a member of aquaporins (A AQP1 is expressed in most tissues i variety of diseases, including cance glycosylated form can also be obser Author Pu Lili Huang 39 Qing Ye 38	n the mammalian body. Alterations r. The predicted molecular weight or ved around 40-45 kDa. (1530176) bmed ID Journal 584501 J Cell Mol Med	of AQP1 expression have been linked to AQP1 is around 28 kDa, while highly Application IF	
Notable Publications	AQP1 is a member of aquaporins (A AQP1 is expressed in most tissues i variety of diseases, including cance glycosylated form can also be obser Author Pu Lili Huang 39 Qing Ye 38	n the mammalian body. Alterations r. The predicted molecular weight of ved around 40-45 kDa. (1530176) bmed ID Journal 584501 J Cell Mol Med 591641 Clin Transl Med 991139 Adv Sci (Weinh)	of AQP1 expression have been linked to AQP1 is around 28 kDa, while highly Application IF IF	
Background Information Notable Publications Storage	AQP1 is a member of aquaporins (A AQP1 is expressed in most tissues i variety of diseases, including cance glycosylated form can also be obser Author Pu Lili Huang 39 Qing Ye 38 Xin Zhang 37 Storage: Store at -20°C. Stable for one year a Storage Buffer:	n the mammalian body. Alterations r. The predicted molecular weight of ved around 40-45 kDa. (1530176) bmed ID Journal 584501 J Cell Mol Med 591641 Clin Transl Med 991139 Adv Sci (Weinh) fter shipment.	of AQP1 expression have been linked to AQP1 is around 28 kDa, while highly Application IF IF	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data








Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 66805-1-lg (AQP1 antibody) at dilution of 1:400 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 66805-1-Ig (AQP1 antibody) at dilution of 1:400 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human kidney tissue using AQP1 antibody (66805-1-Ig, Clone: 2E5D6) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).