

For Research Use Only

CYLD Monoclonal antibody

Catalog Number: 66858-1-Ig



Basic Information

Catalog Number: 66858-1-Ig	GenBank Accession Number: BC012342	Purification Method: Protein A purification
Size: 150UL, Concentration: 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 1540	CloneNo.: 1G2F4
Source: Mouse	Full Name: cylindromatosis (turban tumor syndrome)	Recommended Dilutions: WB 1:500-1:3000
IsoType: IgG2a	Calculated MW: 107 kDa	
Immunogen Catalog Number: AG28333	Observed MW: 110 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB: pig brain tissue, HEK-293 cells, A431 cells
Species Specificity: Human, mouse, rat, pig	

Background Information

CYLD, also named as CYLD1, belongs to the peptidase C67 family. It is the protease that specifically cleaves 'Lys-63'-linked polyubiquitin chains. CYLD has endodeubiquitinase activity and plays an important role in the regulation of pathways leading to NF-kappa-B activation. CYLD contributes to the regulation of cell survival, proliferation and differentiation via its effects on NF-kappa-B activation. It is a negative regulator of Wnt signaling. CYLD inhibits HDAC6 and thereby promotes acetylation of alpha-tubulin and stabilization of microtubules. CYLD plays a role in the regulation of microtubule dynamics, and thereby contributes to the regulation of cell proliferation, cell polarization, cell migration, and angiogenesis. It is required for normal cell cycle progress and normal cytokinesis. CYLD inhibits nuclear translocation of NF-kappa-B and plays a role in the regulation of inflammation and the innate immune response, via its effects on NF-kappa-B activation. It is dispensable for the maturation of intrathymic natural killer cells, but required for the continued survival of immature natural killer cells. CYLD negatively regulates TNFRSF11A signaling and osteoclastogenesis.


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.1% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66858-1-Ig (CYLD antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.