

For Research Use Only

AIM2 Monoclonal antibody

Catalog Number: 66902-1-Ig



Basic Information

Catalog Number: 66902-1-Ig	GenBank Accession Number: BC010940	Purification Method: Protein A purification
Size: 150ul , Concentration: 2000 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 9447	CloneNo.: 3E12A8
Source: Mouse	Full Name: absent in melanoma 2	Recommended Dilutions: WB 1:1000-1:4000
Isotype: IgG1	Calculated MW: 343 aa, 39 kDa	
Immunogen Catalog Number: AG14607	Observed MW: 37-39 and 53 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : RAW 264.7 cells, rabbit spleen tissue, human heart tissue, Sp2/0 cells
Species Specificity: Human, mouse, rat	

Background Information

Absent in melanoma 2 (AIM2) is a sensor of cytosolic DNA that is responsible for activation of the inflammasome and host immune responses to DNA viruses and intracellular bacteria [PMID: 22695634]. Upon binding to DNA, it is thought to undergo oligomerization and to associate with PYCARD initiating the recruitment of caspase-1 precursor and processing of interleukin-1 beta and interleukin-18. Otherwise, it functions as a tumor suppressor which may act by repressing NF-kappa-B transcriptional activity [PMID:16432157,20566831]. 66902-1-Ig antibody detects the 37-39 and 53 kDa protein in SDS-PAGE similar to the paper published. [PMID: 28809949]


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66902-1-Ig (AIM2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.