

For Research Use Only

# ANP32A Monoclonal antibody

Catalog Number: 67687-1-Ig **2 Publications**



## Basic Information

<b>Catalog Number:</b> 67687-1-Ig	<b>GenBank Accession Number:</b> BC007200	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 150ul , Concentration: 1000 ug/ml by Nanodrop and 500 ug/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 8125	<b>CloneNo.:</b> 2D8C5
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> P39687	<b>Recommended Dilutions:</b> WB 1:2000-1:10000 IHC 1:200-1:800 IF/ICC 1:200-1:800
<b>Isotype:</b> IgG2a	<b>Full Name:</b> acidic (leucine-rich) nuclear phosphoprotein 32 family, member A	
<b>Immunogen Catalog Number:</b> AG8546	<b>Calculated MW:</b> 249 aa, 29 kDa	
	<b>Observed MW:</b> 29 kDa	

## Applications

<b>Tested Applications:</b> WB, IHC, IF/ICC, ELISA	<b>Positive Controls:</b>
<b>Cited Applications:</b> WB	<b>WB :</b> A549 cells, HSC-T6 cells, HuH-7 cells, HeLa cells, Raji cells, Jurkat cells, Sp2/0 cells
<b>Species Specificity:</b> human, mouse, rat	<b>IHC :</b> mouse brain tissue, human kidney tissue
<b>Cited Species:</b> human	<b>IF/ICC :</b> A431 cells,
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	

## Background Information

ANP32A, an acidic leucine-rich nuclear phosphoprotein, is a member of the inhibitor of the histone acetyltransferase (INHAT) complex and preferentially binds to unmodified histone H3 tails in vitro. ANP32A is expressed in normal tissues as well as in pancreatic, breast and prostate cancer and other malignant tumors. ANP32A plays an important role in cell proliferation, apoptosis, transcriptional regulation, signal transduction and other processes.

## Notable Publications

Author	Pubmed ID	Journal	Application
Liuke Sun	39737943	Nat Commun	WB
Liuke Sun	37436988	Sci Adv	WB

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol, pH7.3  
**Aliquoting is unnecessary for -20°C storage**

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67687-1-Ig (ANP32A antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 67687-1-Ig (ANP32A antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 67687-1-Ig (ANP32A antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using ANP32A antibody (67687-1-Ig, Clone: 2D8C5 ) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using ANP32A antibody (67687-1-Ig, Clone: 2D8C5 ) at dilution of 1:800 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002), CL594-Phalloidin (red).