

For Research Use Only

EIF3B Monoclonal antibody

Catalog Number: 68202-1-Ig



Basic Information

Catalog Number: 68202-1-Ig	GenBank Accession Number: BC001173	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 8662	CloneNo.: 1G12G12
Source: Mouse	UNIPROT ID: P55884	Recommended Dilutions: WB 1:2000-1:10000
Isotype: IgG1	Full Name: eukaryotic translation initiation factor 3, subunit B	
Immunogen Catalog Number: AG31427	Calculated MW: 93 kDa	
	Observed MW: 115 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HCT 116 cells, A431 cells, HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells
Species Specificity: Human, mouse, rat	

Background Information

EIF3B, also named Eukaryotic translation initiation factor 3 subunit B, is an 814 amino acid protein, which contains 1 RRM (RNA recognition motif) domain and 8 WD repeats and belongs to the eIF-3 subunit B family. EIF3B as an RNA-binding component of the eukaryotic translation initiation factor 3 (eIF-3) complex, is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAⁱ and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation, and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression. The calculated molecular weight of EIF3B is 93 kDa, but the phosphorylated EIF3B protein is about 115 kDa.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68202-1-Ig (EIF3B antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control.