For Research Use Only

Cytokeratin 6A Monoclonal antibody Catalog Number:68294-1-Ig 1 Publications



Basic Information	Catalog Number: 68294-1-lg	GenBank Accession Numl BC008807	ber: Purification Method: Protein G purification
	Size:	GeneID (NCBI):	CloneNo.:
	150ul , Concentration: 1000 ug/ml by Nanodrop and 899 ug/ml by Bradford method using BSA as the standard;		1G1D7
		UNIPROT ID: P02538	Recommended Dilutions: WB 1:5000-1:50000
	Source: Mouse	Full Name: keratin 6A	IHC 1:1150-1:4600
	Isotype: IgG1	Calculated MW: 60 kDa	
	Immunogen Catalog Number: AG0882	Observed MW: 60 kDa	
Applications	Tested Applications: WB, IHC, ELISA	Po	ositive Controls:
	Cited Applications:		B : HaCaT cells, A431 cells, human saliva, rat skin ssue, mouse skin tissue
	IF		IC : human cervical cancer tissue, human oesophag
	Species Specificity: Human, mouse, rat	cancer tissue	
	Cited Species: mouse		
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen	
Background Information	TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0 Keratins are a large family of protein are classified into two major sequence including K9-K23, and the hair keratir	vely, antigen ith citrate s that form the intermedia te types. Type I keratins ar ns Ha 1-Ha 8. Type II keratir he hair keratins, Hb 1-Hb6.	te filament cytoskeleton of epithelial cells, which e a group of acidic intermediate filament proteins, is are the basic or neutral courterparts to the acidic Keratin 6 is a type II keratin. It is used as a marker
Background Information	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0 Keratins are a large family of protein are classified into two major sequence including K9-K23, and the hair keratire type I keratins, including K1-K8, and te for epidermal hyperproliferation and	vely, antigen ith citrate s that form the intermedia te types. Type I keratins ar ns Ha 1-Ha 8. Type II keratir he hair keratins, Hb 1-Hb6.	e a group of acidic intermediate filament proteins, as are the basic or neutral courterparts to the acidic
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0 Keratins are a large family of protein are classified into two major sequence including K9-K23, and the hair kerative type I keratins, including K1-K8, and te for epidermal hyperproliferation and Author Pub	vely, antigen ith citrate s that form the intermedia se types. Type I keratins ar ns Ha1-Ha8. Type II keratir he hair keratins, Hb1-Hb6. differentiation.	e a group of acidic intermediate filament proteins, is are the basic or neutral courterparts to the acidic Keratin 6 is a type II keratin. It is used as a marker Application
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0 Keratins are a large family of protein are classified into two major sequence including K9-K23, and the hair kerative type I keratins, including K1-K8, and te for epidermal hyperproliferation and Author Pub	vely, antigen ith citrate s that form the intermedia se types. Type I keratins ar is Ha1-Ha8. Type II keratir the hair keratins, Hb1-Hb6. differentiation. med ID Journal \$28409 Cancer C er shipment. % glycerol pH 7.3.	e a group of acidic intermediate filament proteins, is are the basic or neutral courterparts to the acidic Keratin 6 is a type II keratin. It is used as a marker Application

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 68294-1-1g (Cytokeratin 6 antibody) at dilution of 1:2300 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 68294-1-Ig (Cytokeratin 6 antibody) at dilution of 1:2300 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Various lysates were subjected to SDS PAGE followed by western blot with 68294-1-1g (Cytokeratin 6 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.