

For Research Use Only

Phospho-IkB Alpha (Ser32/36) Monoclonal antibody

Catalog Number: 68999-1-Ig **1 Publications**



Basic Information

Catalog Number:

68999-1-Ig

Size:

100ul , Concentration: 1000 ug/ml by Nanodrop;

Source:

Mouse

Isotype:

IgG1

GenBank Accession Number:

BC004983

GeneID (NCBI):

4792

UNIPROT ID:

P25963

Full Name:

nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha

Calculated MW:

36 kDa

Observed MW:

36-39 kDa

Purification Method:

Protein G purification

CloneNo.:

2G8E5

Recommended Dilutions:

WB: 1:2000-1:10000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity:

human

Cited Species:

mouse

Positive Controls:

WB : PC-3 cells, MCF-7 cells, Calyculin A treated MCF-7 cells, Calyculin A treated PC cells

Background Information

Nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha (NFKBIA, synonyms: IKBA, MAD-3, NFKB1). NFKB1 or NFKB2 is bound to REL, RELA, or RELB to form the NFKB complex. The NFKB complex is inhibited by I-kappa-B proteins (NFKBIA or NFKBIB), which inactivate NF-kappa-B by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-kappa-B proteins by kinases (IKBKA, or IKBKB) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B complex. Activated NFKB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime. The antibody also detects the phosphorylated form of the NFKBIA (IkB Alpha).

Notable Publications

Author	Pubmed ID	Journal	Application
Qingyue Zhang	40106903	Int Immunopharmacol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Untreated PC-3 cells, PC-3 cells (treated with 100 nM Calyculin A for 0.5 hours) were subjected to SDS PAGE followed by western blot with 68999-1-Ig (Phospho-IkB Alpha antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH (HRP-60004) antibody as loading control.



Untreated MCF-7 cells, MCF-7 cells (treated with 100 nM Calyculin A for 0.5 hours) were subjected to SDS PAGE followed by western blot with 68999-1-Ig (Phospho-IkB Alpha antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH (HRP-60004) antibody as loading control.