

For Research Use Only

NeutraKine® IL-4 Monoclonal antibody



Catalog Number: 69005-1-Ig

Basic Information

Catalog Number: 69005-1-Ig	GenBank Accession Number: GeneID (NCBI): 3565	Purification Method: Protein G purification
Size: 100ug	Full Name: interleukin 4	CloneNo.: 1H5G5
Source: Mouse		
Isotype: IgG1		
Immunogen Catalog Number: HZ-1004		

Applications

Tested Applications:
Neutralization, ELISA

Species Specificity:
Human

Background Information

IL4 is a cytokine produced by CD4+ T cells in response to helminthes and other extracellular parasites. It promotes the proliferation and differentiation of antigen presenting cells. IL4 also plays a pivotal role in antibody isotype switching and stimulates the production of IgE. This cytokine has been applied in the treatment of autoimmune disorder like multiple myeloma, cancer, psoriasis, and arthritis. IL4 has also been extensively applied to inhibit detrimental effect of Th1. It may promote the growth of epithelial tumors by mediating increased proliferation and survival.

This antibody can be used to neutralize the bioactivity of IL-4.

Storage

Storage:
Lyophilized antibodies are stable for 1 year from the date of receipt if stored between (-20°C) and (-80°C). Upon reconstitution we recommend that the solution can be stored at (4°C) for short term or at (-20°C) to (-80°C) for long term. Repeated freeze thaw cycles should be avoided with reconstituted products.

Storage Buffer:
Sterile PBS.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Recombinant human IL-4 (Cat.NO. HZ-1004) stimulates proliferation of TF-1 cell line (human erythroleukemic cell line) in a dose-dependent manner (blue curve, refer to bottom X-left Y). The activity of human IL-4 (0.5 ng/mL HZ-1004) is neutralized by mouse anti-human IL-4 monoclonal antibody 69005-1-Ig at serial dose (red curve, refer to top X-right Y). The ND50 is typically 200-500 ng/mL.