

For Research Use Only

# NeutraKine® IL-17A Mouse McAb

Catalog Number: 69021-1-Ig **3 Publications**



## Basic Information

Catalog Number:	69021-1-Ig	GenBank Accession Number:	Purification Method:
Size:	100ug	GeneID (NCBI):	Protein G purification
Source:	Mouse	3605	CloneNo.:
Isotype:	IgG1	ENSEMBL Gene ID:	1F3E3
Immunogen Catalog Number:	HZ-1113	ENSG00000112115	Recommended Dilutions:
		Full Name:	Neutralization: 1:10-1:100
		interleukin 17A	ELISA: 1:10-1:100

## Applications

Tested Applications:	Positive Controls:
ELISA, Neutralization	Neutralization : HT-1080 cells, HT-1080 cell
Cited Applications:	ELISA : Recombinant protein,
Neutralization, Cell treatment	
Species Specificity:	
Human	
Cited Species:	
human	

## Background Information

IL17A, also named as IL-17, is a proinflammatory cytokine. IL-17, synthesized only by memory T cells and natural killer cells, has pleiotropic effects, mainly in the recruitment and activation of neutrophils. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. The IL-17 receptor is a type I transmembrane protein, that is widely expressed on epithelial cells, fibroblasts, B and T cells, and monocytic cells. In psoriatic skin lesions, both Th17 cells and their downstream effector molecules, e.g. IL-17 and IL-22, are highly increased.

This antibody can be used to neutralize the bioactivity of IL-17A.

## Notable Publications

Author	Pubmed ID	Journal	Application
Danyang Li	36267325	Int J Chron Obstruct Pulmon Dis	Neutralization
Guanzhan Liang	38423357	Cell Mol Gastroenterol Hepatol	Neutralization
Kevin M Motz	37159282	JCI Insight	Cell treatment

## Storage

**Storage:**  
Lyophilized antibodies are stable for 1 year from the date of receipt if stored between (-20°C) and (-80°C). Upon reconstitution we recommend that the solution can be stored at(4°C) for short term or at(-20°C) to (-80°C) for long term. Repeated freeze thaw cycles should be avoided with reconstituted products.

**Storage Buffer:**  
Sterile PBS, pH7.4


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Recombinant human IL-17A (Cat.NO. HZ-1113) stimulates HT-1080 cells (human fibroblast cell line) produce IL-6 in a dose dependent manner (blue curve, refer to bottom X-Left Y axis). The activity of human IL-17A (40ng/mL HZ-1113) is neutralized by mouse anti-human IL-17A monoclonal antibody 69021-1-Ig (red curve, refer to top X-right Y axis). The ND50 is typically 200-600ng/mL.



Recombinant human IL-17A (Cat.NO. HZ-1113) stimulates HT-1080 cells (human fibroblast cell line) produce IL-6 in a dose dependent manner (blue curve, refer to bottom X-Left Y axis). The activity of human IL-17A (40ng/mL HZ-1113) is neutralized by mouse anti-human IL-17A monoclonal antibody 69021-1-Ig (red curve, refer to top X-right Y axis). The ND50 is typically 200-600ng/mL. The NeutraControl mouse anti-human IL-17A monoclonal antibody 69521-



Indirect ELISA was carried out by coating recombinant Human IL-17A (Cat.NO. HZ-1113) at 70 ng/well followed by blocking and adding serial diluted IL-17A antibody 69021-1-Ig and 69521-1-Ig respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.

