

For Research Use Only

CD3 Recombinant antibody

Catalog Number: 81324-1-RR

Featured Product

1 Publications



Basic Information

Catalog Number: 81324-1-RR	GenBank Accession Number: BC049847	Purification Method: Protein A purification
Size: 100ul , Concentration: 500 ug/ml by Nanodrop;	GeneID (NCBI): 916	CloneNo.: 4I19
Source: Rabbit	ENSEMBL Gene ID: ENSG00000198851	Recommended Dilutions: WB: 1:5000-1:50000 IHC: 1:5000-1:20000 IF-P: 1:50-1:500
Isotype: IgG	UNIPROT ID: P07766	
Immunogen Catalog Number: AG11797	Full Name: CD3e molecule, epsilon (CD3-TCR complex)	
	Calculated MW: 207 aa, 23 kDa	
	Observed MW: 23-25 kDa	

Applications

Tested Applications: WB, IHC, IF-P, ELISA	Positive Controls:
Cited Applications: IHC	WB : Jurkat cells, MOLT-4 cells, pig thymus tissue, rat thymus tissue, mouse thymus tissue
Species Specificity: human, mouse, rat, pig	IHC : human colon tissue, human tonsillitis tissue, mouse small intestine tissue, mouse spleen tissue
Cited Species: human	IF-P : mouse spleen tissue, mouse small intestine tissue
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

CD3 is a complex of proteins that directly associates with the T cell receptor (TCR). The TCR/CD3 complex of T-lymphocytes consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta. The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction. CD3 is considered to be a pan-T cell marker for detection of normal and neoplastic T cells. This antibody is directed against the epsilon chain of human CD3 molecule.

Notable Publications

Author	Pubmed ID	Journal	Application
Enkui Zhang	40065365	J Transl Med	IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of CD3 antibody (81324-1-RR; 1:60000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CD3 transfected Jurkat cells.




Various lysates were subjected to SDS PAGE followed by western blot with 81324-1-RR (CD3E antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 81324-1-RR (CD3 antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 81324-1-RR (CD3 antibody) at dilution of 1:10000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 81324-1-RR (CD3 antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse small intestine tissue using CD3 antibody (81324-1-RR, Clone: 4I19) at dilution of 1:1000 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse spleen tissue using CD3 antibody (81324-1-RR, Clone: 4I19) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 81324-1-RR against Human CD3 were performed. The affinity constant is 1.13 nM.