

For Research Use Only

IBA1 Recombinant antibody

Catalog Number: 81728-1-RR **5 Publications**



Basic Information

Catalog Number: 81728-1-RR	GenBank Accession Number: BC009474	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 199	CloneNo.: 4D5
Source: Rabbit	ENSEMBL Gene ID: ENSG00000204472	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000 IF-P 1:250-1:1000 IF-Fro 1:200-1:800
Isotype: IgG	UNIPROT ID: P55008	
Immunogen Catalog Number: AG1363	Full Name: allograft inflammatory factor 1	
	Calculated MW: 17 kDa	
	Observed MW: 17 kDa	

Applications

Tested Applications: WB, IHC, IF-P, FC (Intra), ELISA	Positive Controls: WB : THP-1 cells, U-937 cells, HL-60 cells
Cited Applications: IHC, IF	IHC : rat brain tissue, human liver tissue, mouse brain tissue, mouse liver tissue
Species Specificity: human, mouse, rat	IF-P : rat brain tissue,
Cited Species: mouse, rat	IF-Fro : mouse brain tissue,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

IBA1 is a 143 amino acid cytoplasmic, inflammation response scaffold protein. It is constitutively expressed in monocytes and macrophages and is known to be involved in macrophage activation. It is a marker of activated macrophage. Expression of IBA1 is up-regulated in activated microglia following facial nerve axotomy, ischemia, and several brain diseases, thereby implicating it in the activated phenotypes of microglia.

Notable Publications

Author	Pubmed ID	Journal	Application
Juan Gao	39654367	CNS Neurosci Ther	IF
Rui Wang	39600046	Adv Mater	IHC
Wei Shen	39489209	Exp Cell Res	IF

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




THP-1 cells were subjected to SDS PAGE followed by western blot with 81728-1-RR (IBA1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 81728-1-RR (IBA1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 81728-1-RR (IBA1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using IBA1 antibody (81728-1-RR, Clone: 4D5) at dilution of 1:500 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




Various lysates were subjected to SDS PAGE followed by western blot with 81728-1-RR (IBA1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using IBA1 antibody (81728-1-RR, Clone: 4D5) at dilution of 1:500 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat brain tissue using IBA1 antibody (81728-1-RR, Clone: 4D5) at dilution of 1:250 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse brain tissue using IBA1 antibody (81728-1-RR, Clone: 4D5) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1×10^6 THP-1 cells were intracellularly stained with 0.4 ug IBA1 Recombinant antibody (81728-1-RR, Clone:4D5) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2) (red), or 0.4 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).