For Research Use Only

ATG5 Recombinant antibody

Catalog Number:81803-1-RR Featured Product



Basic Information

Catalog Number: GenBank Accession Number:

81803-1-RR BC002699 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 9474 5M5 Nanodrop: **UNIPROT ID:**

Q9H1Y0 WB 1:5000-1:50000 Rabbit

Full Name: Isotype: ATG5 autophagy related 5 homolog

(S. cerevisiae) IgG

Immunogen Catalog Number: Calculated MW:

AG0214 32 kDa

Observed MW: 50-55 kDa

Applications

Tested Applications:

WB, IHC, IP, ELISA Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method: Protein A purification

CloneNo.:

Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:500-1:2000

Positive Controls:

WB: HeLa cells, Jurkat cells, HepG2 cells, A549 cells, NIH/3T3 cells, RAW 264.7 cells, HSC-T6 cells, PC-12

cells

IP: HeLa cells,

IHC: human ovary tumor tissue,

Background Information

ATG5, also named as APG5L and ASP, belongs to the ATG5 family. It is required for autophagy. It plays an important role in the apoptotic process, possibly within the modified cytoskeleton. Its expression is a relatively late event in the apoptotic process, occurring downstream of caspase activity. Autophagy is a catabolic process for the autophagosomic-lysosomal degradation of bulk cytoplasmic contents. Formation of the autophagosome involves a ubiquitin-like conjugation system in which Atg12 is covalently bound to Atg5 and targeted to autophagosome vesicles. It mediates autophagosome-independent host protection. This antibody is raised against 28-275 amino acids of human ATG5. It can recognize the ATG5-ATG12 complex (55 kDa) which can be truncated and generate a 40-45 kDa band.

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 81803-1-RR (ATG5 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




WB result of ATG5 antibody (81803-1-RR; 1:6000; room temperature for 1.5 hours) with wild-type and ATG5 knockout HeLa cells.



IP result of anti-ATG5 (IP:81803-1-RR, 4ug; Detection:81803-1-RR 1:2000) with HeLa cells lysate 1280 ug.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 81803-1-RR (ATG5 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 81803-1-RR against Human ATG5 were performed. The affinity constant is 42.4 nM.