For Research Use Only

IGF2BP3 Recombinant antibody

Catalog Number:81805-1-RR 1 Publications



Basic Information

Catalog Number: GenBank Accession Number: 81805-1-RR BC065269

81805-1-RR BC065269 Protein A purification
Size: GenelD (NCBI): CloneNo.:
100ul , Concentration: 1000 ug/ml by 10643 1L11

Full Name:

Nanodrop; UNIPROT ID: Recommended Dilutions: Source: 000425 WB 1:5000-1:50000

Isotype: insulin-like growth factor 2 mRNA

IgG binding protein 3

Immunogen Catalog Number: Calculated MW: AG6226 64 kDa

Observed MW: 64 kDa IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:200-1:800

Purification Method:

Applications

Tested Applications: WB, IHC, IP, ELISA

Cited Applications:

IHC

Rabbit

Species Specificity: human, mouse, rat Cited Species: Positive Controls:

WB: HEK-293 cells, HepG2 cells, HeLa cells, K-562 cells, NIH/3T3 cells, HSC-T6 cells, mouse placenta

IP: HeLa cells,

r . HeLa cetts,

IHC: human pancreas cancer tissue, human cervical squamous cancer tissue, human colon cancer tissue

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

IGF2BP3, also named IMP3, KOC1, and VICKZ3, belongs to the RRM IMP/VICKZ family. It is one of the RNA binding proteins involved in mRNA localization and translational control. IGF2BP3 is expressed during embryogenesis, as well as in some malignant tumors. It can be used as an independent prognostic factor for osteosarcoma. Both isoforms (64kd and 22kd) of IGFBP3 can be recognized by this antibody. And IGFBP3 is nuclear and cytoplasm stains.

Notable Publications

Author	Pubmed ID	Journal	Application
Li Yang	38714753	Sci Rep	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 81805-1-RR (IGF 2BP3 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




IP result of anti-IGF2BP3 (IP:81805-1-RR, 4ug; Detection:81805-1-RR 1:20000) with HeLa cells lysate 1280 ug.




IP result of anti-IGF2BP3 (IP:81805-1-RR, 4ug; Detection:81805-1-RR 1:4000) with HeLa cells lysate 1280 ug.




Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 81805-1-RR (IGF2BP3 antibody) at dilution of 1:400 (under 0x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 81805-1-RR (IGF2BP3 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human cervical squamous cancer tissue slide using 81805-1-RR (IGF2BP3 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 81805-1-RR (IGF2BP3 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 81805-1-RR (IGF2BP3 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 81805-1-RR (IGF2BP3 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 81805-1-RR against Human I GF2BP3 were performed. The affinity constant is below 1 pM.