

For Research Use Only

# APC Anti-Mouse IL-4 (11B11)

Catalog Number: APC-65074



## Basic Information

Catalog Number:

APC-65074

Size:

100ug, 0.2 mg/ml

Source:

Rat

Isotype:

IgG1, kappa

GenBank Accession Number:

BC027514

GeneID (NCBI):

16189

UNIPROT ID:

P07750

Full Name:

interleukin 4

Purification Method:

Affinity purification

CloneNo.:

11B11

Excitation/Emission maxima  
wavelengths:

650 nm / 660 nm

## Applications

Tested Applications:

FC (Intra)

Species Specificity:

mouse

## Background Information

Interleukin-4 (IL-4), a member of the  $\alpha$ -helical cytokine family, is produced by activated CD4<sup>+</sup> T cells, basophils, and mast cells. It promotes the proliferation and differentiation of antigen presenting cells. IL-4 also plays a pivotal role in antibody isotype switching and stimulates the production of IgE. This cytokine has been applied in the treatment of autoimmune disorder like multiple myeloma, cancer, psoriasis, and arthritis. IL-4 has also been extensively applied to inhibit detrimental effect of Th1. It may promote the growth of epithelial tumors by mediating increased proliferation and survival. (PMID: 24489573;3049907;21663408)

## Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide and 0.5% BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10<sup>6</sup> unstimulated or PMA and ionomycin stimulated (in the presence of protein transport inhibitors) mouse Th2-polarized splenocytes were surface stained with 0.2 ug PE Anti-Mouse CD4 (GK1.5) (PE-65104, Clone: GK1.5). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C), and then intracellularly stained with 0.5 ug APC Anti-Mouse IL-4 (APC-65074, Clone:11B11).