## For Research Use Only

## APC Anti-Human CD28 (CD28.2)

Catalog Number: APC-65099 1 Publications



**Basic Information** 

Catalog Number:

APC-65099

Size:

100 tests , 5  $\mu$ l/test

Source: Mouse

Isotype: IgG1, kappa ENSG00000178562 UNIPROT ID: P10747 Full Name:

> CD28 molecule Calculated MW:

GenBank Accession Number:

220 aa, 25 kDa

BC093698

940

GeneID (NCBI):

ENSEMBL Gene ID:

**Purification Method:** 

Affinity purification

CloneNo.: CD28.2

Recommended Dilutions:

FC: 5 ul per  $10^6$  cells in  $100 \mu l$ 

suspension

Excitation/Emission maxima

wavelengths: 650 nm / 660 nm

**Applications** 

**Tested Applications:** 

FC

Cited Applications:

FC (Intra)

Species Specificity:

Human **Cited Species:** 

human

Positive Controls:

FC: human PBMCs,

**Background Information** 

CD28 (T-cell-specific surface glycoprotein CD28), also known as T44 and Tp44, is a 44 kD disulfide-linked homodimeric type I glycoprotein (PMID: 2162180). It is a member of the immunoglobulin superfamily and is expressed on most T lineage cells, NK cell subsets, and plasma cells (PMID: 2162180, 8386518). CD28 may affect in vivo immune responses by functioning both as a cell adhesion molecule linking B and T lymphocytes and as the surface component of a novel signal transduction pathway (PMID: 2162180, 3021470). CD28 binds both CD80 and CD86 with a highly conserved motif MYPPY in the CDR3-like loop (PMID: 15696168, 7964482). CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death (PMID: 1348520).

**Notable Publications** 


Author **Pubmed ID** Application Journal 39382426 Cardiovasc Res FC (Intra) Xiang Li

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

## Selected Validation Data



1X10^6 human PBMCs were surface co-stained with FITC Anti-Human CD3 and 5 ul APC Anti-Human CD28 (APC-65099, Clone:CD28.2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul APC Anti-Human CD28 (APC-65099, Clone:CD28.2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.