## For Research Use Only

## APC Anti-Human PD-1/CD279 (J110)

Catalog Number: APC-65119



**Basic Information** 

Catalog Number:

APC-65119

Size:

100tests , 5 µl/test

Source: Mouse

Isotype:

lgG1

GenBank Accession Number:

BC074740 GeneID (NCBI):

**UNIPROT ID:** 

Q15116 Full Name:

programmed cell death 1

Calculated MW:

288 aa, 32 kDa

**Purification Method:** 

Affinity purification

CloneNo.: J110

Recommended Dilutions:

FC: 5 ul per  $10^6$  cells in  $100 \mu l$ 

suspension

Excitation/Emission maxima

wavelengths: 650 nm / 660 nm

**Applications** 

**Tested Applications:** 

Species Specificity:

human

Positive Controls:

FC: human peripheral blood lymphocytes,

## **Background Information**

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.1% gelatin, pH7.2

## **Selected Validation Data**



1X10^6 PHA-stimulated (5 µg/mL, overnight) human peripheral blood lymphocytes were surface stained with 5.00 ul APC Anti-Human PD-1/CD279 (APC-65119, Clone: J110) and FITC Anti-Human CD3 (FITC-65151, Clone: UCHT1). Cells were not fixed.



1X10^6 human peripheral blood lymphocytes were surface stained with 5.00 ul APC Anti-Human PD-1/CD279 (APC-65119, Clone: J110) and FITC Anti-Human CD3 (FITC-65151, Clone: UCHT1). Cells were not fixed.