For Research Use Only

APC Anti-Human CD73 (AD2)

Catalog Number: APC-65162



Basic Information

Catalog Number: APC-65162

BC015940 GeneID (NCBI):

Size: 100 tests , 5 μ l/test 4907

UNIPROT ID: Source: Mouse P21589 Isotype: Full Name:

IgG1, kappa 5'-nucleotidase, ecto (CD73)

> Calculated MW: 29 kDa, 63 kDa

GenBank Accession Number:

Applications

Tested Applications: FC

Species Specificity:

Human

Purification Method: Affinity purification

CloneNo.: AD2

Recommended Dilutions:

FC: 5 ul per 10^6 cells in 100 μ l

suspension

Positive Controls:

FC: human PBMCs,

Excitation/Emission maxima

wavelengths: 650 nm / 660 nm

Background Information


CD73, also known as ecto-5'-nucleotidase (5'-NT), is a 70-kDa, glycosyl-phosphatidylinositol-linked membranebound glycoprotein found in most tissues (PMID: 18404475; 20179192). CD73 is an ectoenzyme that catalyzes the dephosphorylation of AMP and other nucleoside monophosphates (PMID: 9553767). In the human immune system, CD73 is expressed on subsets of T and B cells, on germinal center follicular dendritic cells, and on thymic medullary reticular fibroblasts and epithelial cells (PMID: 2137649; 9553767). CD73 is highly expressed in many human solid tumors and is closely involved in cancer progression (PMID: 20179192).

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



1X10^6 human PBMCs were surface co-stained with CoraLite® Plus 405 CD3 and 5 ul APC Anti-Human CD73 (APC-65162, Clone:AD2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul APC Anti-Human CD73 (APC-65162, Clone:AD2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.