For Research Use Only

Biotin-conjugated PARK7/DJ-1 Polyclonal antibody

Catalog Number:Biotin-11681

Featured Product




| Basic Information | Catalog Number: Biotin-11681 | GenBank Accession Number: BC008188 | Purification Method: Antigen affinity purification | | |
|--|---|---|---|---|--|
| | Size: 100ul , Concentration: 1000 µg/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG2287 | UNIPROT ID: Q99497 Full Name: Parkinson disease (autosomal | Recommended Dilutions: IHC 1:400-1:1600 | | |
| | | | | recessive, early onset) 7 Calculated MW: 189 aa, 20 kDa | |
| | | | | Observed MW: 20 kDa, 25 kDa | |
| | | Applications | Tested Applications: IHC | Positive Controls: | |
| | | | IHC : mouse kidney tissue, Species Specificity: human, mouse, rat | | |
| Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 | | | | | |
| Background Information | PARK7, also named as DJ1, belongs to the peptidase C56 family. It protects cells against oxidative stress and cell death. PARK7 plays a role in regulating expression or stability of the mitochondrial uncoupling proteins SLC25A14 and SLC25A27 in dopaminergic neurons of the substantia nigra pars compacta and attenuates the oxidative stress induced by calcium entry into the neurons via L-type channels during pacemaking. It eliminates hydrogen peroxide and protects cells against hydrogen peroxide-induced cell death. PARK7 has cell-growth promoting activity and transforming activity. It may function as a redox-sensitive chaperone. It's precursor undergoes a cleavage of a C-terminal peptide and subsequent activation of protease activity in response to oxidative stress. The amino acid replace at 166 (L \rightarrow P) reduces PARK7 protein stability and leads to increased degradation. The predicted MW of this protein is 20 kDa, An additional 25 kDa band can be observed due to modification (PMID: 31767755). | | | | |
| | Storage: Store at -20°C. Avoid exposure to ligh | It. Stable for one year after shipme | nt. | | |
| Storage | Storage Buffer: PBS with 50% Glycerol, 0.05% Proclir | 1300, 0.5% BSA, pH 7.3. | | | |

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using Biotin-11681 (PARK7/DJ-1 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using Biotin-11681 (PARK7/DJ-1 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).