For Research Use Only

CoraLite® Plus 488-conjugated ubiquitin Polyclonal antibody

Catalog Number: CL488-10201



Purification Method:

IF-P 1:50-1:500

wavelengths:

493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Basic Information

Applications

Catalog Number: GenBank Accession Number:

CL488-10201 BC000379
Size: GeneID (NCBI):

100ul , Concentration: 1000 ug/ml by 7314
Nanodrop; UNIPROT ID:
Source: POCG47

Rabbit Full Name:
Isotype: ubiquitin B

IgG

Immunogen Catalog Number:

AG0260

Tested Applications: Positive Controls:

IF-P, FC (Intra)

IF-P: human pancreas cancer tissue,

Species Specificity: human, mouse, rat

Background Information

Ubiquitin B (UBB) is a member of ubiquitin family, one of the most conserved proteins known. Ubiquitin B is required for ATP-dependent, non-lysosomal intracellular protein degradation of abnormal proteins and normal proteins with a rapid turnover. Ubiquitin B is covalently bound to proteins to be degraded, and presumably labels these proteins for degradation. Ubiquitin also binds to histone H2A in actively transcribed regions but does not cause histone H2A degradation, suggesting that ubiquitin is also involved in regulation of gene expression. When polyubiquitin is free (unanchored-polyubiquitin), it also has distinct roles, such as in activation of protein kinases, and in signaling. This gene consists of three direct repeats of the ubiquitin coding sequence with no spacer sequence. Consequently, the protein is expressed as a polyubiquitin precursor with a final amino acid after the last repeat. Aberrant form of this protein has been noticed in patients with Alzheimer's and Down syndrome. Interestingly ubiquitin also becomes covalently bonded to many types of pathological inclusions which appear to be resistant to normal degradation.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human pancreas cancer tissue using CoraLite® Plus 488 ubiquitin antibody (CL488-10201) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human ubiquitin (CL488-10201) (red), or 0.8 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).