For Research Use Only

CoraLite® Plus 488-conjugated TBP Monoclonal antibody

www.ptglab.com

Catalog Number: CL488-66166

Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** CL488-66166 BC110341 Protein A purification

> GeneID (NCBI): CloneNo.: 100ul, Concentration: 1000 ug/ml by 6908 2H3B2

Nanodrop: **UNIPROT ID:** Recommended Dilutions:

P20226 IF/ICC: 1:50-1:500 Mouse

FC (Intra): 0.20 ug per 10^6 cells in a Full Name: 100 µl suspension

Isotype: TATA box binding protein Excitation/Emission maxima IgG2a Calculated MW:

wavelengths: Immunogen Catalog Number: 338 aa, 38 kDa 493 nm / 522 nm

AG12383

Applications Tested Applications: Positive Controls:

IF/ICC, FC (Intra) IF/ICC: C2C12 cells, A431 cells

Species Specificity: FC (Intra): HeLa cells, human, mouse, rat, pig

Background Information

The TATA binding protein (TBP) is a transcription factor that binds specifically to a DNA sequence TATA box. This DNA sequence is found about 25-30 base pairs upstream of the transcription start site in some eukaryotic gene promoters. TBP, along with a variety of TBP-associated factors, make up the TFIID, a general transcription factor that in turn makes up part of the RNA polymerase II preinitiation complex. As one of the few proteins in the preinitation complex that binds DNA in a sequence-specific manner, it helps position RNA polymerase II over the transcription start site of the gene. However, it is estimated that only 10-20% of human promoters have TATA boxes. Therefore, TBP is probably not the only protein involved in positioning RNA polymerase II. This antibody detects human TBP (~40 kDa) and mouse/rat Tbp (~35 kDa).

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed C2C12 cells using CL488-66166 (TBP antibody) at dilution of 1:50.



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite® Plus 488 Anti-Human TBP (CL488-66166, Clone:2H3B2) (red), or 0.2 ug Mouse IgG2a Isotype Control (CL488-66360-2, Clone: K11A1B2A2) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using CoraLite® Plus 488 TBP antibody (CL488-66166, Clone: 2H3B2) at dilution of 1:100.