For Research Use Only

CoraLite® Plus 488-conjugated Caspase 9/P35/P10 Monoclonal antibody



Catalog Number: CL488-66169

Basic Information

Catalog Number: GenBank Accession Number: Purification Method: Protein A purification

 Size:
 GeneID (NCBI):
 CloneNo.:

 100ul , Concentration: 1000 ug/ml by 842
 1B7G2

Nanodrop; UNIPROT ID: Recommended Dilutions:
Source: P55211 IF/ICC 1:50-1:500

Mouse Full Name: Excitation/Emission maxima

 Isotype:
 caspase 9, apoptosis-related cysteine wavelengths:

 IgG2b
 peptidase

 493 nm / 522 nm

Immunogen Catalog Number: Calculated MW: AG20813 46 kDa

Observed MW: 46 kDa, 35 kDa

Applications

Tested Applications:

IF/ICC, FC (Intra)

Positive Controls:

IF/ICC : HeLa cells,

Species Specificity: human, mouse

Background Information

Caspase 9, apoptosis-related cysteine protease (CASP9,synonyms: MCH6, APAF3, APAF-3, ICE-LAP6, CASPASE-9c)is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme. Capase 9 is processed by caspase APAF1; this step is thought to be one of the earliest in the caspase activation cascade.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using Coralite® Plus 488 Caspase 9/P35/P10 antibody (CL488-66169, Clone: 1B7G2) at dilution of 1:200.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Coralite® Plus 488 Anti-Human Caspase 9/P35/P10 (CL488-66169, Clone:1B7G2) (red), or 0.4 ug Mouse IgG2b Isotype Control (CL488-66360-3, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).