

For Research Use Only

CoraLite® Plus 488-conjugated c-Fos Monoclonal antibody

Catalog Number: CL488-66590

Featured Product



Basic Information

Catalog Number:

CL488-66590

Size:

100ul , Concentration: 1000 ug/ml by Nanodrop;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG24340

GenBank Accession Number:

BC004490

GeneID (NCBI):

2353

UNIPROT ID:

P01100

Full Name:

FOS

Calculated MW:

41 kDa

Observed MW:

55-60 kDa

Purification Method:

Protein G purification

CloneNo.:

1G2C5

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

human, mouse, rat

Background Information

c-FOS, also named as Proto-oncogene c-Fos and G0/G1 switch regulatory protein 7, is a 380 amino acid protein, which contains 1 bZIP (basic-leucine zipper) domain and belongs to the bZIP family. FOS is expressed at very low levels in quiescent cells. When cells are stimulated to reenter growth, FOS undergo 2 waves of expression, the first one peaks 7.5 minutes following FBS induction. At this stage, the FOS protein is localized endoplasmic reticulum. The second wave of expression occurs at about 20 minutes after induction and peaks at 1 hour. At this stage, the FOS protein becomes nuclear. The calculated molecular weight of FOS is 40 kDa, but Phosphorylated FOS protein is about 60-65 kDa. It is involved in important cellular events, including cell proliferation, differentiation and survival; genes associated with hypoxia; and angiogenesis; which makes its dysregulation an important factor for cancer development. It can also induce a loss of cell polarity and epithelial-mesenchymal transition, leading to invasive and metastatic growth in mammary epithelial cells. Expression of c-fos is an indirect marker of neuronal activity because c-fos is often expressed when neurons fire action potentials. Upregulation of c-fos mRNA in a neuron indicates recent activity

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ HeLa cells were intracellularly stained with 0.8 ug CoraLite® Plus 488-conjugated c-Fos Monoclonal antibody (CL488-66590, Clone:1G2C5) (red), or 0.8 ug CoraLite® Plus 488 Mouse IgG1 Isotype Control (MOPC-21) (CL488-65124, Clone: MOPC-21) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).