For Research Use Only

## CoraLite® Plus 488-conjugated FBXO6 Monoclonal antibody



**Purification Method:** 

CloneNo.: 3B7F9

wavelengths:

493 nm / 522 nm

Protein A purification

**Recommended Dilutions:** IF/ICC 1:50-1:500

Excitation/Emission maxima

Catalog Number: CL488-67697

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-67697 BC020880 GeneID (NCBI):

Size: 100ul, Concentration: 1000 ug/ml by 26270

Nanodrop; **UNIPROT ID:** 

Q9NRD1 Source Mouse Full Name:

Isotype: F-box protein 6 lgG2b Calculated MW:

Immunogen Catalog Number: 293 aa, 34 kDa AG30512 Observed MW:

34 kDa

**Applications** 

**Tested Applications:** 

IF/ICC

Species Specificity: human, mouse, rat

Positive Controls:

IF/ICC: HEK-293 cells,

## **Background Information**

FBXO6(F-box only protein 6) is also named as FBG2, FBS2, FBX6 and belongs to the F-box protein family. FBXO6 is the substrate recognition component of a Skp1-Cullin1-F-box protein (SCF) ubiquitin E3 ligase complex, recognizing the chitobiose in unfolded N-glycoprotein to target glycoproteins for polyubiquitination and degradation. It can regulate Chk1 ubiquitination and degradation in both normally cycling cells and during replication stress. This protein is mainly detected in the cytoplasm in both cultured cells and in tumor tissues. Its expression is a predictive biomarker of tumor responsiveness to the important anticancer drugs. Positive Fbxo6 expression correlated with early TNM stage and favorable prognosis of NSCLC patients. (PMID:19716789, PMID: 27855403, PMID: 31140586).


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using Coralite® Plus 488 FBXO6 antibody (CL488-67697, Clone: 3B7F9) at dilution of 1:200.



1X10^6 Jurkat cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human FBXO6 (CL488-67697, Clone:3B7F9) (red), or 0.4 ug Mouse IgG2b Isotype Control (CL488-66360-3, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).