For Research Use Only

CoraLite® Plus 488-conjugated AGO2 Monoclonal antibody



Catalog Number: CL488-67934

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL488-67934 BC007633 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 27161 Nanodrop: **UNIPROT ID:** Q9UKV8 Mouse Full Name:

Isotype: eukaryotic translation initiation

factor 2C, 2 lgG1 Immunogen Catalog Number: Calculated MW: AG27772 97 kDa

Observed MW: 90 kDa

Applications

Tested Applications: IF/ICC, FC (Intra)

Species Specificity: human, mouse, rat

Purification Method:

Protein G purification

CloneNo.: 1G2H12

Recommended Dilutions: IF/ICC 1:500-1:2000

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

Background Information

EIF2C2, also named as AGO2, belongs to the argonaute family and Ago subfamily. It is required for RNA-mediated gene silencing (RNAi) by the RNA-induced silencing complex (RISC). The 'minimal RISC' appears to include EIF2C2/AGO2 bound to a short guide RNA such as a microRNA (miRNA) or short interfering RNA (siRNA). EIF2C2 may inhibit translation initiation by binding to the 7-methylguanosine cap, thereby preventing the recruitment of the translation initiation factor eIF4-E. It also inhibit translation initiation via interaction with EIF6, which itself binds to the 60S ribosomal subunit and prevents its association with the 40S ribosomal subunit. EIF2C2 can also upregulate the translation of specific mRNAs under certain growth conditions.

Positive Controls:

IF/ICC: HeLa cells,

Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Coralite® Plus 488 AGO2 antibody (CL488-67934, Clone: 1G2H12) at dilution of 1:1000.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Coralite® Plus 488 Anti-Human AGO2 (CL488-67934, Clone:1G2H12) (red), or 0.4 ug x. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).