

For Research Use Only

# CoraLite®555-conjugated TDRKH Monoclonal antibody

Catalog Number:CL555-66845



## Basic Information

<b>Catalog Number:</b> CL555-66845	<b>GenBank Accession Number:</b> BC032690	<b>Purification Method:</b> Protein G purification
<b>Size:</b> 100ul , Concentration: 1000 ug/ml by Nanodrop;	<b>GeneID (NCBI):</b> 11022	<b>CloneNo.:</b> 2B1B12
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> Q9Y2W6	<b>Recommended Dilutions:</b> IF/ICC 1:50-1:500
<b>Isotype:</b> IgG1	<b>Full Name:</b> tudor and KH domain containing	<b>Excitation/Emission maxima wavelengths:</b> 557 nm / 570 nm
<b>Immunogen Catalog Number:</b> AG4720	<b>Calculated MW:</b> 606 aa, 67 kDa	
	<b>Observed MW:</b> 67-70 kDa	

## Applications

<b>Tested Applications:</b> IF/ICC, FC (Intra)	<b>Positive Controls:</b> IF/ICC : U2OS cells,
<b>Species Specificity:</b> human	

## Background Information

Tudor and KH domain-containing protein(TDRKH)Tudor domains are protein modules that mediate protein-protein interactions, potentially by binding to methylated ligands. Both KH and tudor domains are involved in binding RNA or single-strand DNA. Tudor also is a germ cell-specific protein with multiple Tudor domains and is involved in germ plasm formation and germ cell specification. TDRKH is one Tudor protein with a single Tudor domain.


## Storage

**Storage:**  
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.  
**Aliquoting is unnecessary for -20°C storage**


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



$1 \times 10^6$  MCF-7 cells were intracellularly stained with 0.4  $\mu$ g CoraLite®555 Anti-Human TDRKH (CL555-66845, Clone:2B1B12) (red), or 0.4  $\mu$ g Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using CoraLite®555 TDRKH antibody (CL555-66845, Clone: 2B1B12 ) at dilution of 1:200.