For Research Use Only

## CoraLite®555-conjugated SUMO2/3 Monoclonal antibody

www.ptglab.com

Catalog Number: CL555-67154

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL555-67154 BC008450 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 6613 Nanodrop: **UNIPROT ID:** P61956 Mouse Full Name:

Isotype: SMT3 suppressor of mif two 3lgG2b homolog 2 (S. cerevisiae)

Immunogen Catalog Number: Calculated MW: AG28672 11 kDa

> Observed MW: 18 kDa

**Purification Method:** 

Protein A purification

CloneNo.: 1A1B3

Excitation/Emission maxima

wavelengths: 557 nm / 570 nm

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity: Human, Mouse, Rat

## **Background Information**

Ubiquitin is most famous for its function in targeting proteins for degradation by the 26S proteasome, ubiquitin needs to be attached to a substrate in chains (polyubiquitylation) before being recognized by proteasome. Similarly, SUMO (small ubiquitin-related modifier) can be linked to substrates in chains (polysumoylation), SUMO modification has been implicated in many important cellular processes including the control of genome stability, signal transduction, targeting to and formation of nuclear compartments, cell cycle and meiosis. There are 4 confirmed SUMO isoforms in human, SUMO-1, SUMO-2, SUMO-3 and SUMO-4. SUMO-2 and SUMO-3 are nearly  $identical\ but\ are\ distinct\ from\ SUMO-1.\ SUMO2/3\ conjugation\ was\ recently\ widely\ involved\ in\ neuroprotective$ activities. A substitution (M55V) of SUMO 4 was strongly associated with the pathogenesis of type 1 diabetes (T1D) involving NF kappa B related mechanisms.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



1X10^6 Jurkat cells were intracellularly stained with 0.4 ug Coralite®555 Anti-Human SUMO2/3 (CL555-67154, Clone:1A1B3) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).