

For Research Use Only

# CoraLite®555-conjugated c-MYC Monoclonal antibody

Catalog Number: CL555-67447

Featured Product



## Basic Information

Catalog Number:

CL555-67447

Size:

100ul , Concentration: 1000 ug/ml by Nanodrop;

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG17519

GenBank Accession Number:

BC000141

GeneID (NCBI):

4609

UNIPROT ID:

P01106

Full Name:

v-myc myelocytomatosis viral oncogene homolog (avian)

Calculated MW:

49 kDa

Observed MW:

55-60 kDa

Purification Method:

Protein G purification

CloneNo.:

3D9C12

Excitation/Emission maxima wavelengths:

557 nm / 570 nm

## Applications

Tested Applications:

FC (Intra)

Species Specificity:

Human, mouse, rat

## Background Information

MYC contains one basic helix-loop-helix (bHLH) domain. This protein is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. It seems to activate the transcription of growth-related genes. It binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma.

## Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10<sup>6</sup> K-562 cells were intracellularly stained with 0.4 ug CoraLite®555 Anti-Human c-MYC (CL555-67447, Clone:3D9C12) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).