For Research Use Only

CoraLite®555-conjugated RPA2 Monoclonal antibody

Catalog Number: CL555-67999



Basic Information

Catalog Number: GenBank Accession Number:

CL555-67999 BC001630 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 6118 Nanodrop: **UNIPROT ID:** P15927 Mouse Full Name:

Isotype: replication protein A2, 32kDa

lgG1 Calculated MW:

Immunogen Catalog Number: 29 kDa AG24783

Observed MW:

30 kDa

1B9H11 Recommended Dilutions:

Purification Method:

Protein G purification

CloneNo.:

IF/ICC 1:500-1:2000

Excitation/Emission maxima wavelengths: 557 nm / 570 nm

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity:

human, mouse, rat, pig

Positive Controls:

IF/ICC: A431 cells,

Background Information

RPA2, a subunit of the heterotrimeric replication protein A (RPA) complex, is overexpressed in various cancers. RPA2 and the replication protein A (RPA) complex, is overexpressed in various cancers. RPA2 and RPA2 are replication protein A (RPA) complex, is overexpressed in various cancers. RPA2 and RPA2 are replication protein A (RPA) complex, is overexpressed in various cancers. RPA2 are replication protein A (RPA) complex, is overexpressed in various cancers. RPA2 are replication protein A (RPA) complex are replication and replication protein A (RPA) complex are replication and replication are replicated and replication are replicated are replicated are replicated are replicated and replication are replicated are reis a major target for upstream kinases, including CDKs and the three PIKKs (ATR, ataxia-telangiectasia mutated [ATM], and DNA-PK). RPA2 comprises multiple critical Ser/Thr residues that are phosphorylated in sequential order in response to genotoxic stress.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed A431 cells using CoraLite®555 RPA2 antibody (CL555-67999, Clone: 1B9H11) at dilution of 1:1000.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite®555 Anti-Human RPA2 (CL555-67999, Clone:1B9H11) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).