For Research Use Only

CoraLite®555-conjugated Phospho-Hamartin/TSC1 (Ser511) Recombinant antibody




Catalog Number:CL555-80340

Catalog Number: CL555-80340	GenBank Accession Number: NM_000368	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by	GeneID (NCBI): iml by 7248 UNIPROT ID: Q92574 Full Name: tuberous sclerosis 1 Calculated MW: 130 kDa Observed MW:	CloneNo.: 3N6 Excitation/Emission maxima wavelengths: 557 nm / 570 nm
Nanodrop; Source: Rabbit Isotype: IgG		
Tested Applications: FC (Intra) Species Specificity: human, mouse		
mental retardation, autism, and tumo (TSC1)/TSC2 tumor suppressor compl complex function may contribute to to and in vivo, and phosphorylation at th induces mTOR activation. (PMID: 1769	rs in the brain, retina, kidney, hear ex serves as a repressor of the mT umorigenesis. TSC1 is phosphoryla nese sites enhances dissociation of 93255). TSC1 is also threonine-pho	t, and skin. The tuberous sclerosis 1 DR pathway, and disruption of TSC1/TSC2 ated by IKKb at Ser487 and Ser511 in vitro the TSC1/TSC2 complex, which in turn isphorylated at three sites (Thr417, Ser584,
Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin	1300, 0.5% BSA, pH 7.3.	nt.
	CL555-80340 Size: 100ul, Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Tested Applications: FC (Intra) Species Specificity: human, mouse TSC 1, also named as Hamartin, is a tu mental retardation, autism, and tumo (TSC 1)/TSC 2 tumor suppressor compl complex function may contribute to tu and in vivo, and phosphorylation at tf induces mTOR activation. (PMID: 176 and Thr1047) in a reaction catalyzed I Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin	CL555-80340 NM_000368 Size: Genel D (NCBI): 100ul, Concentration: 1000 ug/ml by 7248 Nanodrop; UNIPROT ID: Source: Q92574 Rabbit Full Name: Isotype: tuberous sclerosis 1 IgG Calculated MW: 130 kDa Observed MW: 130 kDa Observed MW: 150 kDa Species Specificity: human, mouse TSC 1, also named as Hamartin, is a tumor suppressor gene syndrome wimental retardation, autism, and tumors in the brain, retina, kidney, hear (TSC 1)/TSC 2 tumor suppressor complex serves as a repressor of the mTC complex function may contribute to tumorigenesis. TSC 1 is phosphorylation at these sites enhances dissociation of induces mTOR activation. (PMID: 17693255). TSC 1 is also threonine-phc and Thr1047) in a reaction catalyzed by cyclin-dependent kinase 1 (CDW Storage: Storage: Storage: Storage

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 HeLa cells untreated (dashed lines) or treated with Calyculin A (red) were intracellularly stained with 0.5 ug CoraLite®555 Anti-Human Phospho-Hamartin/TSC1 (Ser511) (CL555-80340, Clone:3N6) (red), or 0.5 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with 50% MeOH.