For Research Use Only

CoraLite®594-conjugated PhosphomTOR (Ser2448) Recombinant antibody

Catalog Number: CL594-80596



Basic Information

Catalog Number: GenBank Accession Number: Purification Method: CL594-80596 BC117166 Protein A purification

Size:GeneID (NCBI):CloneNo.:100ul , Concentration: 1000 ug/ml by 24753L18

Nanodrop; Full Name: Recommended Dilutions:

Source: FK506 binding protein 12-rapamycin IF/ICC: 1:50-1:500

Rabbit associated protein 1 FC (Intra): 0.13 ug per 10^6 cells in a

Isotype: Calculated MW: 100 µl suspension

IgG 289 kDa Excitation/Emission maxima

Observed MW: wavelengths: 588 nm / 604 nm

Applications

Tested Applications: Positive Controls:

IF/ICC, FC (Intra)

IE/ICC - DMA troat

IF/ICC : PMA treated HEK-293 cells,
Species Specificity:

56 (Intra) Columbia A treated Held

human, rat FC (Intra): Calyculin A treated HeLa cells, HeLa cells

Background Information

MTOR, also named as FRAP1, FRAP, FRAP2 and RAPT1, belongs to the PI3/PI4-kinase family. MTOR is a Ser/Thr protein kinase that functions as an ATP and amino acid sensor to balance nutrient availability and cell growth. MTOR is kinase subunit of both mTORC1 and mTORC2, which regulate cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino-acids. mTORC2 is also activated by growth factors, but seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTOR is phosphorylated at Ser2448 via the PI3 kinase/Akt signaling pathway and autophosphorylated at Ser2481. mTOR plays a key role in cell growth and homeostasis and may be abnormally regulated in tumors.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 HeLa cells untreated (dashed lines) or treated with Calyculin A (red) were intracellularly stained with 0.13 ug CoraLite®594 Anti-Human Phospho-mTOR (Ser2448) (CL594-80596, Clone:3L18) (red), or 0.13 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with 80% MeOH.



Immunofluorescent analysis of (-20°C Ethanol) fixed PMA treated HEK-293 cells using CoraLite® 594 Phospho-mTOR (Ser2448) antibody (CL594-80596, Clone: 3L18) at dilution of 1:200.