For Research Use Only

CoraLite® Plus 647-conjugated Caspase 9/P35/P10 Polyclonal antibody



Purification Method:

wavelengths: 654 nm / 674 nm

Antigen affinity purification

Excitation/Emission maxima

Catalog Number: CL647-10380

Featured Product

Basic Information

Catalog Number: CL647-10380

GenBank Accession Number: BC002452

GeneID (NCBI):

100ul , Concentration: 1000 $\mu g/ml$ by 842

Nanodrop: **UNIPROT ID:**

P55211 Rabbit Full Name:

Isotype caspase 9, apoptosis-related cysteine

IgG peptidase Immunogen Catalog Number: Calculated MW: AG0404 46 kDa

> Observed MW: 46 kDa, 35 kDa

Applications

Tested Applications:

FC (Intra)

Species Specificity:

human, mouse, rat

Background Information

Caspase 9, apoptosis-related cysteine protease (CASP9, synonyms: MCH6, APAF3, APAF-3, ICE-LAP6, CASPASE-9c)is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme. Caspase 9 is processed by APAF1; this step is thought to be one of the earliest in the caspase activation cascade. 10380-1-AP can recognize the pre- and cleaved- caspase 9. In recent years, the localization of caspase9 was a focus of interest. Beside its cytoplasmic distribution, a very extensive localization study was done on rat brain tissue, where caspase9 was found located predominantly in the nucleus and to a lesser extend in the cytoplasm [PMID: 15541731].

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 HepG2 cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human Caspase 9/P35/P10 (CL647-10380) (red), or 0.2 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).