For Research Use Only

CoraLite® Plus 647-conjugated iNOS Polyclonal antibody

www.ptglab.com

Catalog Number: CL647-18985

Featured Product

1 Publications

Basic Information

Catalog Number: CL647-18985

GenBank Accession Number:

NM 000625 GeneID (NCBI):

100ul , Concentration: 1000 $\mu g/ml$ by 4843

Nanodrop: **UNIPROT ID:** Source: P35228

Rabbit Full Name:

Isotype: nitric oxide synthase 2, inducible IgG

Calculated MW:

131 kDa Observed MW:

110-130 kDa, 65-70 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions: IF/ICC: 1:50-1:500

FC (Intra): 0.20 ug per 10⁶ cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse, rat

Cited Species: mouse

Positive Controls:

IF/ICC: HepG2 cells, FC (Intra): HepG2 cells,

Background Information

NOS2, also named as iNOS and NOS2A, produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. NO is a reactive free radical which acts as a biologic mediator in several processes, including neurotransmission, antimicrobial and antitumoral activities. NOS2 is a nitric oxide synthase which is expressed in liver and is inducible by a combination of lipopolysaccharide and certain cytokines. iNOS has a very short half-life due to rapid degradation by calpain. iNOS monomer is a direct substrate of calpain I and can be cleaved by calpain I at the canonical CaM-binding site(503-532aa) of iNOS, and then a ~70kD band can be detected by western (PMID:11786228). This antibody is specific to NOS2.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|------------|-----------|------------|-------------|
| Fengan Cao | 39806403 | Respir Res | |

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 HepG2 cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human iNOS (CL647-18985) (red), or 0.2 ug isotype control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using CoraLite® Plus 647 iNOS antibody (CL647-18985) at dilution of 1:100.