For Research Use Only

CoraLite® Plus 647-conjugated Galectin-3 Monoclonal antibody



Catalog Number: CL647-60207

Featured Product

Basic Information

Catalog Number: GenBank Accession Number: **Purification Method:** CL647-60207 BC001120 Protein A purification

GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000 $\mu g/ml$ by 3958 1C1B2

Nanodrop: **UNIPROT ID:** Recommended Dilutions: P17931 IF/ICC 1:50-1:500

Mouse Excitation/Emission maxima Full Name:

Isotype: lectin, galactoside-binding, soluble, 3 wavelengths: 654 nm / 674 nm lgG2b

Calculated MW: Immunogen Catalog Number: 26 kDa

AG6753 Observed MW:

31 kDa

Applications

Positive Controls: **Tested Applications:** IF/ICC, FC (Intra) IF/ICC: HeLa cells,

Species Specificity: human, mouse, rat

Background Information

Galectins are a family of animal lectins defined by shared characteristic amino-acid sequences and affinity for β galactose-containing oligosac-charides (PMID: 8063692). Galectin-3, a member of the β -galactoside-binding proteins, contains one carbohydrate recognition domain (CRD) and a proline- and glycine-rich N-terminal domain through which is able to form oligomers (PMID: 14758078). Galectin-3 is widely expressed in many normal tissues and a variety of tumors. It is found intracellularly in nucleus and cytoplasm or secreted outside of cell, being present on the cell surface or in the extracellular space (PMID: 16478649). Galectin-3 is involved in various biological processes including cell growth, adhesion, differentiation, apoptosis, angiogenesis, immune response, neoplastic transformation and metastasis (PMID: 16478649; 14758078).


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CoraLite® Plus 647 Galectin-3 antibody (CL647-60207, Clone: 1C1B2) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.8 ug CoraLite® Plus 647 Anti-Human Galectin-3 (CL647-60207, Clone:1C1B2) (red), or 0.8 ug CoraLite® Plus 647 Mouse IgG2b Isotype Control (MPC-11) (CL647-65128, Clone: MPC-11) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).