

CoraLite® Plus 647-conjugated U2AF35 Monoclonal antibody

Catalog Number: **CL647-60289**

Basic Information

Catalog Number: CL647-60289	GenBank Accession Number: BC001923	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/mL by Nanodrop;	GeneID (NCBI): 7307	CloneNo.: 2B7F11
Source: Mouse	UNIPROT ID: Q01081	Excitation/Emission maxima wavelengths: 654 nm / 674 nm
Isotype: IgG2c	Full Name: U2 small nuclear RNA auxiliary factor 1	
Immunogen Catalog Number: AG0399	Calculated MW: 28 kDa	
	Observed MW: 35-40 kDa	

Applications

Tested Applications:
FC (Intra)

Species Specificity:
human, mouse, rat

Background Information

U2 auxiliary factor (U2AF), comprising a large and a small subunit, is a non-snRNP protein required for the binding of U2 snRNP to the pre-mRNA branch site. U2 (RNU2) small nuclear RNA auxiliary factor 1 (U2AF1, synonyms: RN, FP793, U2AF35, U2AFBP, RNU2AF1) is the small subunit which plays a critical role in both constitutive and enhancer-dependent RNA splicing by directly mediating interactions between the large subunit and proteins bound to the enhancers. The U2AF1 gene is localized to chromosome 21q22.3, which is the critical region for three diseases, progressive myoclonus epilepsy, autoimmune polyglandular disease type 1, and one form of bipolar affective disorder. The calculated molecular weight of U2AF35 is 28 kDa, but the modified U2AF35 protein is about 35-40 kDa.

Storage

Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ HeLa cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human U2AF35 (CL647-60289, Clone:2B7F11) (red), or 0.2 ug APC-65128 ; APC Mouse IgG2b Isotype Control (APC-65128, Clone: MPC-11) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).