For Research Use Only

CoraLite® Plus 647 Anti-Rat CD8a (OX-

www.ptglab.com

Catalog Number: CL647-65180

Basic Information

Catalog Number:

CL647-65180

100ug , 500 μ g/ml

Source: Mouse

Isotype: IgG1, kappa GenBank Accession Number:

BC088126 GeneID (NCBI):

24930

UNIPROT ID: P07725

Full Name: CD8a molecule **Purification Method:**

Purified by protein-A affinity

chromatography

CloneNo.: **0X-8**

Recommended Dilutions:

FC: 0.5 ug per 10^6 cells in $100 \mu l$

suspension

Excitation/Emission maxima wavelengths:

654 nm / 674 nm

Applications

Tested Applications:

Species Specificity:

Positive Controls:

FC: Wistar rat splenocytes,

Background Information


CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8αβ heterodimer while CD8αα homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) $\gamma\delta$ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).

Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide, pH7.3

Selected Validation Data



1X10^6 Wistar rat splenocytes were surface stained with 0.5 ug CoraLite® Plus 647 Anti-Rat CD8a (CL647-65180, Clone: OX-8) (red), or 0.5 ug Control Antibody. Cells were not fixed.