For Research Use Only

## CoraLite® Plus 647-conjugated CDK1 Monoclonal antibody

Catalog Number:

CL647-67575



**Purification Method:** 

Catalog Number: CL647-67575

**Basic Information** 

GenBank Accession Number:

BC014563 Protein A purification

GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000  $\mu g/ml$  by 983 2A11E4

Nanodrop: **UNIPROT ID:** Excitation/Emission maxima

P06493 wavelengths: 654 nm / 674 nm Mouse Full Name:

Isotype: cell division cycle 2, G1 to S and G2 to

lgG2a

Immunogen Catalog Number: Calculated MW: AG29018 25 kDa

> Observed MW: 34 kDa

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity: Human, Mouse, Rat

**Background Information** 

CDK1, also named as CDC2, belongs to the protein kinase superfamily, CMGC Ser/Thr protein kinase family and CDC2/CDKX subfamily. CDK1 plays a key role in the control of the eukaryotic cell cycle. It is required in higher cells for entry into S-phase and mitosis. CDK1 is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. It is a component of the kinase complex that phosphorylates the repetitive C-terminus of RNA polymerase II. Mitotic cyclins stably associate with CDK1 and function as regulatory subunits. CDK1 has 2 isoforms produced by alternative splicing with the molecular mass of 34 kDa and 27 kDa.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human CDK1 (CL647-67575, Clone:2A11E4) (red), or 0.2 ug isotype control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).