

For Research Use Only

CoraLite®647-conjugated TGF beta 1 Monoclonal antibody



Catalog Number:CL647-69012

Basic Information

Catalog Number: CL647-69012	GenBank Accession Number: 7040	Purification Method: Protein G purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 7040	CloneNo.: 1E3E9
Source: Mouse	Full Name: transforming growth factor, beta 1	Excitation/Emission maxima wavelengths: 650 nm/665 nm
Isotype: IgG1		
Immunogen Catalog Number: HZ-1011		

Applications

Tested Applications:
FC (Intra)

Species Specificity:
Human

Background Information

TGFB, also named as LAP and TGFB1, is a multifunctional peptide that controls proliferation, differentiation, and other functions in many cell types. TGFB acts synergistically with TGFA in inducing transformation. It also acts as a negative autocrine growth factor. Dysregulation of TGFB activation and signaling may result in apoptosis. Many cells synthesize TGFB and almost all of them have specific receptors for it. TGFB positively and negatively regulates many other growth factors. It plays an important role in bone remodeling as it is a potent stimulator of osteoblastic bone formation, causing chemotaxis, proliferation and differentiation in committed osteoblasts. It is highly expressed in bone. Mutation of TGFB are the cause of Camurati-Engelmann disease (CED) which known as progressive diaphyseal dysplasia 1 (DPD1).

This antibody is used to Block/Neutralize TGF beta 1 protein.

Storage

Storage:
Store at -20°C. Avoid exposure to light.


Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ HEK-293 cells were intracellularly stained with 0.4 ug CoraLite®647 Anti-Human TGF beta 1 (CL647-69012, Clone:1E3E9) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).