For Research Use Only

CoraLite® Plus 750 Anti-Human CD7 (4H9)

www.ptglab.com

Catalog Number: CL750-65203

Basic Information

Catalog Number: CL750-65203

100 tests , 5 μ l/test

Source: Mouse

IgG2a, kappa

Isotype:

Calculated MW:

GenBank Accession Number:

BC009293 GeneID (NCBI):

ENSEMBL Gene ID: ENSG00000173762 UNIPROT ID:

P09564 Full Name: CD7 molecule

240 aa, 25 kDa

Purification Method:

Affinity purification

CloneNo.: 4H9

Recommended Dilutions:

FC: 5 ul per 10^6 cells in $100 \mu l$

suspension

Excitation/Emission maxima

wavelengths: 755 nm / 780 nm

Applications

Tested Applications:

Species Specificity:

human

Positive Controls:

FC: human PBMCs,

Background Information

CD7, also known as Leu-9 or GP40, is a 40-kDa single-pass type I transmembrane glycoprotein belonging to the immunoglobulin superfamily. CD7 is expressed on thymocytes, T cells, NK cells, and cells in the early stages of T, B, and myeloid cell differentiation (PMID: 10530432). It is one of the earliest antigens to appear on cells of the Tlymphocyte lineage (PMID: 3501369). CD7 plays a significant role in T-cell and T-cell/B-cell interactions during early lymphoid development. It is involved in the regulation of cell signaling and biology, including T-cell activation, proliferation, and the expression of interleukin-2 receptor alpha (IL-2Ra) (PMID: 11485208). CD7 is not only a marker for T-cells but also has functional implications in immune cell interactions and adhesion. It can provide co-stimulatory signals and is associated with CD3 and CD45, participating in T-cell activation (PMID: 7523512).

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 human PBMCs were surface co-stained with APC Anti-Human CD4 and 5 ul Coralite® Plus 750 Anti-Human CD7 (CL750-65203, Clone:4H9) or Mouse IgG2a Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul CoraLite® Plus 750 Anti-Human CD7 (CL750-65203, Clone:4H9) or Mouse IgG2a Isotype Control. Cells were not fixed. Lymphocytes were gated.