

For Research Use Only

# HRP-conjugated PCNA Monoclonal antibody

Catalog Number:HRP-60097

Featured Product

2 Publications



## Basic Information

<b>Catalog Number:</b> HRP-60097	<b>GenBank Accession Number:</b> BC000491	<b>Purification Method:</b> Protein G purification
<b>Size:</b> 100ul , Concentration: 1000 ug/ml by Nanodrop;	<b>GeneID (NCBI):</b> 5111	<b>CloneNo.:</b> 10D10E11
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> P12004	<b>Recommended Dilutions:</b> WB 1:5000-1:50000
<b>Isotype:</b> IgG1	<b>Full Name:</b> proliferating cell nuclear antigen	
<b>Immunogen Catalog Number:</b> AG7416	<b>Calculated MW:</b> 29 kDa/31 kDa	
	<b>Observed MW:</b> 36-38 kDa	

## Applications

<b>Tested Applications:</b> WB	<b>Positive Controls:</b> WB : HEK-293 cells,
<b>Cited Applications:</b> WB	
<b>Species Specificity:</b> human, mouse, pig, rat	
<b>Cited Species:</b> human, mouse	

## Background Information

Proliferating Cell Nuclear Antigen, commonly known as PCNA, is a protein that acts as a processivity factor for DNA polymerase  $\delta$  in eukaryotic cells. This protein is an auxiliary protein of DNA polymerase delta and is involved in the control of eukaryotic DNA replication by increasing the polymerase's processibility during elongation of the leading strand. PCNA induces a robust stimulatory effect on the 3'-5' exonuclease and 3'-phosphodiesterase, but not apurinic-apyrimidinic (AP) endonuclease, APEX2 activities. It has to be loaded onto DNA in order to be able to stimulate APEX2. PCNA protein is highly conserved during evolution; the deduced amino acid sequences of rat and human differ by only 4 of 261 amino acids. PCNA has been used as loading control for proliferating cells. The calculated molecular weight of PCNA is 29 kDa, but modified PCNA is 36kDa (PMID: 1358458).

## Notable Publications

Author	Pubmed ID	Journal	Application
Ha T Nguyen	36468795	Mol. Reprod Dev	WB
Shengjie Zhu	39638770	Phytother Res	WB


## Storage

**Storage:**  
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with HRP-60097 (PCNA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.