

## HumanKine® recombinant human IL-17F protein



Animal Component-Free

Human cell expressed

Tag-Free

Endotoxin Free

### Product Description

Animal-free Recombinant Human IL-17F is a homodimeric glycoprotein with an apparent molecular mass of 40 kDa. This cytokine is produced in a human cell expression system with serum-free, chemically defined media. Production in human 293 cells offers authentic glycosylation, which contributes to stability in cell growth media and other applications. IL-17 is commonly associated with allergic responses. IL-17 induces the production of many other cytokines (such as IL-6, G-CSF, GM-CSF, IL-1beta, TGF beta, and TNF alpha), chemokines (including IL-8, GRO alpha, and MCP-1), and prostaglandins (e.g., PGE2) from many cell types (fibroblasts, endothelial cell, epithelial cells, keratinocytes, and macrophages). The release of cytokines causes many functions, such as airway remodeling, a characteristic of IL-17 responses.

**Alternative Names** Cytokine ML 1, IL 17F, IL17, IL-17, IL17F, IL-17F, interleukin 17F, ML 1, ML1


**Source** Human Embryonic Kidney cells (HEK293). HEK293-derived IL-17F protein

**Species Reactivity** human


### Specifications

Test	Method	Specification
Activity (EC50)	Dose-dependent induction of alkaline phosphatase production in a HEK293 reporter cell line.	4.5-22.5 ng/mL
Molecular Mass	SDS-PAGE	22 kDa reduced, 40 kDa non-reduced, homodimer, glycosylated
Purity	SDS-PAGE	>95%
Endotoxin	LAL	<1 EU/μg

### Activity Data



### SDS-PAGE



Preparation	
Shipping Temperature	ambient temperature
Formulation	1x PBS, See Certificate of Analysis for details
Reconstitution	Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein to 0.2 mg/mL in sterile 1x PBS pH 7.4 containing 0.1% endotoxin-free recombinant human serum albumin (HSA). Gently swirl or tap vial to mix.

Stability and Storage	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
	Lyophilized	Room Temperature	2 weeks
	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
Avoid repeated freeze-thaw cycles.			

[www.ptglab.com](http://www.ptglab.com)

Document #: FR-QA118-101 Rev 0  
Data Sheet Version #: 1

**Proteintech Group, Inc.**  
5500 Pearl Street, Suite 400 Rosemont, IL 60612  
t: 1-888-478-4522; f: 1-312-455-8408  
Email: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)