

Catalog Number: HZ-1192

Data Sheet





Animal Component-Free

Human cell expressed

Tag-Free


Endotoxin Free

Product Description


Animal-free Recombinant Human M-CSF is expressed in human 293 cells and has an apparent molecular mass of 35 to 40 kDa due to glycosylation. M-CSF is a potent hematopoietic factor produced by a variety of cells including lymphocytes, monocytes, fibroblasts, endothelial cells, myoblasts, and osteoblasts. It is a key regulator of cellular proliferation, differentiation, and survivial of blood monocytes, tissue macrophages, and their progenitor cells. This cytokine is produced in a human cell expression system with serum-free, chemically defined media.

Alternative Names	ative Names CSF 1, CSF1, Lanimostim, M CSF, MCSF, M-CSF	
Source	Human Embryonic Kidney cells (HEK293). HEK293-derived M-CSF protein	
Species Reactivity	human,mouse	

Specifications Specification Specification Specification Specification Specification Specification Specificatio					
Test	Method	Specification			
Activity (EC50)	Dose-dependent stimulation of the proliferation of murine M-NFS-60 cells (Mouse myeloid leukemia indicator cell line)	0.7-4.0 ng/mL			
Molecular Mass	SDS-PAGE	19 and 23 kDa reduced, 35 and 40 kDa non-reduced, homodimer, glycosylated			
Purity	SDS-PAGE	>95%			
Endotoxin	LAL	<1 EU/µg			



Recombinant human M-CSF (Cat no: HZ-1192) stimulates dose-dependent proliferation of the murine mouse myloid leukemia (M-NFS-60) cell line. Cell number was quantitatively assessed by Prestoblue® Cell Viability Reagent. M-NFS-60 cells were treated with increasing concentrations of recombinant M-CSF for 48 hours. The EC50 range is 0.7-4.0 ng/mL.



www.ptglab.com

Document #: FR-QA118-101 Rev 0
Data Sheet Version #: 1

Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com

	Preparation				
Shipping Temperature	ambient temperature				
Formulation 1x PBS, See Certificate of Analysis for details					
Reconstitution	Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein to 0.2 mg/mL in sterile 1x PBS pH 7.4 containing 0.1% endotoxin-free recombinant human serum albumin (HSA). Gently swirl or tap vial to mix.				

Stability and Storage	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
	Lyophilized	Room Temperature	2 weeks
	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
		Avoid repeated freeze-thaw cycles.	

www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com