

Recombinant Human PDGFbb

Catalog Number: **HZ-1308**

HEK293 expressed

Endotoxin-free

Animal-component free

Technical Specifications

Species	human
Expression	HEK293
Activity	Typically ≤ 5 ng/mL EC50
Purity	>95%
Endotoxin	<1 EU/ μ g
Molecular Mass	29 to 32 kDa, homodimer, glycosylated
Formulation	50mM Acetate pH 3.5 + 500mM NaCl, See Certificate of Analysis for details
Gene ID	5155

Reconstitution Buffer

Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in sterile 10 mM HOAc containing 0.1% endotoxin-free recombinant human serum albumin (HSA).

Stability and Storage


Lyophilized proteins are stable for 1 year from the date of receipt if stored between (-20°C) and (-80°C). Upon reconstitution we recommend that the solution can be stored at (4°C) for short term or at (-20°C) to (-80°C) for long term. Repeated freeze thaw cycles should be avoided with reconstituted products.

Product Description


Animal-free Recombinant Human PDGFbb is a growth factor that promotes blood vessel formation, mitogenesis, chemotaxis, etc. The PDGF family members (four homodimers and one heterodimer) are secreted, disulfide-linked dimeric glycoproteins that regulate their cellular functions through interaction with PDGFR receptors. PDGFbb is synthesized, stored, and released by alpha granules of platelets. It is also known as PDGF-2, becaplermin, or GDGF. Dysregulation in PDGF signaling has been shown to be associated with tumorigenesis and progression of cancer. Recombinant PDGFbb is used in treatment of chronic ulcers and to speed healing in surgical procedures.

Synonyms

Becaplermin, c sis, FLJ12858, PDGF 2, PDGF subunit B, PDGF2, PDGFB, PDGFbb, Proto oncogene c Sis, SIS, SSV



Animal-free Recombinant human PDGFbb (Catalogue: HZ-1308) stimulates dose-dependent proliferation of the NIH/3T3 mouse fibroblast cell line. Viable cell number was quantitatively assessed by Prestobule Cell Viability Reagent. NIH/3T3 cells were plated and serum starved with 0.1% FBS for 24 hours and added increasing concentrations of recombinant human PDGFbb for 48 hours before addition of Prestobule Cell Viability Reagent. EC50 is less than 5 ng/mL as determined by a 4 parameter non-linear regression model.



The protein was resolved by SDS-polyacrylamide gel electrophoresis and the gel was stained with Coomassie blue.

For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or
1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

Humankine[®] product line
HUMANZYME
Now part of Proteintech Group[®]