

For Research Use Only

PE Anti-Human CD154 (24-31)

Catalog Number: PE-65156



Basic Information

Catalog Number:
PE-65156

Size:
100tests, 10 µl/test

Source:
Mouse

Isotype:
Mouse IgG1

GenBank Accession Number:
BC071754

GeneID (NCBI):
959

Full Name:
CD40 ligand

Calculated MW:
261 aa, 29 kDa

Purification Method:

The purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography.

CloneNo.:
24-31

Excitation/Emission maxima wavelengths:
496 nm, 565 nm / 578 nm

Applications

Tested Applications:
FC

Species Specificity:
Human, Non-human primates

Background Information

The CD40 ligand (CD40L, TRAP, CD154), a member of the TNF superfamily of ligands, is expressed as either a 33-kd transmembrane homologue or 18-kd soluble form (sCD154). CD40L is primarily expressed on activated CD4+ T cells and on a small proportion of CD8+ T cells and platelets. It binds to CD40 on antigen-presenting cells (APC), which leads to many effects depending on the target cell type. It has been suggested that CD40/CD40L interactions regulate oxidative stress and affect various signaling pathways in both the immunological and the cardiovascular systems. The CD40/CD40L system is also involved in tumorigenesis. Its expression is tightly regulated, and abnormal levels of CD40L are associated with the pathogenesis of atheromatous plaque destabilization and thrombotic events. Multiple mutations in CD40LG gene have been identified that are associated with hyper-IgM immunodeficiency syndrome type 1.

Storage

Storage:
Store at 2-8°C. Avoid exposure to light.
Storage Buffer:
PBS with 0.1% sodium azide.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1×10^6 PMA (50 ng/ml) and ionomycin (1 ug/ml)-stimulated human PBMCs (5 hours) were surface stained with 10 ul PE Anti-Human CD154 (PE-65156, Clone: 24-31) (red) or isotype control antibody (black). Cells were not fixed.