

RAB6B Polyklonaler Antikörper

Katalog-Nr.: 10340-1-AP

3 Publikationen

Allgemeine Informationen

Katalog-Nr.:	GenBank-Zugangsnummer:	Reinigungsmethode:
10340-1-AP	BC002510	Antigen-Affinitätsreinigung
Größe:	GenID (NCBI):	Empfohlene Verdünnungen:
150ul, Konzentration: 450 µg/ml von Nanodrop und 233 µg/ml durch die Bradford-Methode mit BSA als Standard;	51560	WB 1:500-1:2000 IP 0.5-4.0 ug für IP und 1:500-1:2000 für WB IHC 1:20-1:200 IF 1:10-1:100
Wirt:	Vollständiger Name:	
Kaninchen	RAB6B, member RAS oncogene family	
Isotyp:	Berechneté Masse:	
IgG	23 kDa	
Immunogen Katalognummer:	Beobachteté Masse:	
AG0322	24 kDa	

Anwendungen

Geprüfte Anwendungen:	Positivkontrollen:
IF, IHC, IP, WB, ELISA	WB: Maushirngewebe, C6-Zellen, Rattenhirngewebe
In Publikationen genannte Anwendungen:	IP: Maushirngewebe,
IF, WB	IHC: humanes Gliomgewebe,
Getestete Reaktivität:	IF: C6-Zellen,
Human, Maus, Ratte	
Zitierte Arten:	
Maus, Ratte	
Hinweis-IHC: Antigendemaskierung mit TE-Puffer pH 9,0 empfohlen. (*) Wahlweise kann die Antigendemaskierung auch mit Citratpuffer pH 6,0 erfolgen.	

Hintergrundinformationen

The human RAB genes share structural and biochemical properties with the Ras gene superfamily. Accumulating data suggests an important role for RAB proteins either in endocytosis or in biosynthetic protein transport. The transport of newly synthesized proteins from endoplasmic reticulum to the Golgi complex and to secretory vesicles involves the movement of carrier vesicles, a process that appears to involve RAB protein function. Rab6A has been shown to be a regulator of membrane traffic from the Golgi apparatus towards the endoplasmic reticulum (ER). Rab6B is encoded by an independent gene which is located on chromosome 3 region q21-q23. In contrast to Rab6A whose expression is ubiquitous, Rab6B is expressed in a tissue and cell-type specific manner. Rab6B is predominantly expressed in brain and the neuroblastoma cells. In brain, Rab6B was found to be specifically expressed in microglia, pericytes and Purkinje cells. Endogenous Rab6B localises to the Golgi apparatus and to ERGIC-53-positive vesicles. Comparable studies between Rab6A and Rab6B revealed distinct biochemical and cellular properties. Rab6B displays lower GTP-binding activities and is distributed over Golgi and ER membranes, whereas Rab6A is more restricted to the Golgi apparatus. Since the GTP-bound form of Rab6B does interact with all known Rab6A effectors, including Rabkinesin-6, the results suggest a cell-type specific role for Rab6B in retrograde membrane traffic at the level of the Golgi complex.

Bemerkenswerte Veröffentlichungen

Verfasser	Pubmed ID	Journal	Anwendung
Elisa Ghelfi	29760588	Proteome Sci	WB
Liyuan Guo	34332492	J Neuroimmunol	IF
Sabine Bardin	35979738	EMBO Rep	WB

Lagerung




Lagerungsbedingungen:
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil
Lagerungspuffer:
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


Ausgewählte Validierungsdaten



mouse brain tissue were subjected to SDS PAGE followed by western blot with 10340-1-AP (RAB6B antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP Result of anti-RAB6B (IP:10340-1-AP, 3ug; Detection:10340-1-AP 1:1000) with mouse brain tissue lysate 4000ug.



Immunofluorescent analysis of C6 cells using 10340-1-AP (RAB6B antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).