

Nur für Forschungszwecke

c-SRC Polyklonaler Antikörper

Katalog-Nr.:11097-1-AP

60 Publikationen



Allgemeine Informationen

Katalog-Nr.: 11097-1-AP	GenBank-Zugangsnummer: BC011566	Reinigungsmethode: Antigen-Affinitätsreinigung
Größe: 150ul , Konzentration: 800 µg/ml von Nanodrop;	GeneID (NCBI): 6714	Empfohlene Verdünnungen: WB 1:300-1:800 IP 0.5-4.0 ug für IP und 1:500-1:2000 für WB
Wirt: Kaninchen	Vollständiger Name: v-src sarcoma (Schmidt-Ruppin A-2) viral oncogene homolog (avian)	IHC 1:100-1:400 IF 1:10-1:100
Isotyp: IgG	Berechnete Masse: 60 kDa	
Immunogen Katalognummer: AG1575	Beobachtete Masse: 55 kDa, 60 kDa	

Anwendungen

Geprüfte Anwendungen:
FC, IF, IHC, IP, WB, ELISA

In Publikationen genannte Anwendungen:
IF, IHC, IP, WB

Getestete Reaktivität:
Hamster, Human, Maus, Ratte

Zitierte Arten:
Human, Maus, Ratte

Hinweis-IHC: Antigenmaskierung mit TE-Puffer pH 9,0 empfohlen. (*) Wahlweise kann die Antigenmaskierung auch mit Citratpuffer pH 6,0 erfolgen.

Positivkontrollen:

WB: Jurkat-Zellen, A431-Zellen, CHO-Zellen, COLO 320-Zellen, Maushirngewebe, SH-SY5Y-Zellen

IP: SH-SY5Y-Zellen,

IHC: humanes Mammakarzinomgewebe, humanes Hirngewebe, humanes Milzgewebe

IF: A431-Zellen,

Hintergrundinformationen

SRC, also named as SRC1 and p60-Src, belongs to the protein kinase superfamily, Tyr protein kinase family and SRC subfamily. It is a non-receptor protein tyrosine kinase that plays pivotal roles in numerous cellular processes such as proliferation, migration, and transformation. In concert with PTK2B, SRC plays an important role in osteoclastic bone resorption. It promotes energy production in osteoclasts by activating mitochondrial cytochrome C oxidase. SRC can be detected as 60 kDa and 55 kDa (PMID: 15047935, 21461171).

Bemerkenswerte Veröffentlichungen

Verfasser	Pubmed ID	Journal	Anwendung
YanHua Fan	36174847	Fitoterapia	WB
Lianhong Yin	29033837	Front Pharmacol	WB
Jiao Wu	34517345	Aging (Albany NY)	WB

Lagerung

Lagerungsbedingungen:

Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil

Lagerungspuffer:

PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Ausgewählte Validierungsdaten




Jurkat cells were subjected to SDS PAGE followed by western blot with 11097-1-AP (SRC antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




IP Result of anti-SRC (IP:11097-1-AP, 5ug; Detection:11097-1-AP 1:1000) with SH-SY5Y cells lysate 2900ug.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 11097-1-AP (c-SRC Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 11097-1-AP (c-SRC Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of A431 cells using 11097-1-AP (SRC antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10⁶ SH-SY5Y cells were stained with 0.2ug c-SRC antibody (11097-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.