

## Allgemeine Informationen

<b>Katalog-Nr.:</b> 12582-1-AP	<b>GenBank-Zugangsnummer:</b> BC013694	<b>Reinigungsmethode:</b> Antigen-Affinitätsreinigung
<b>Größe:</b> 150ul, Konzentration: 350 µg/ml von Nanodrop;	<b>GeneID (NCBI):</b> 5725	<b>Empfohlene Verdünnungen:</b> WB 1:2000-1:16000 IHC 1:50-1:500 IF 1:50-1:500
<b>Wirt:</b> Kaninchen	<b>Vollständiger Name:</b> polypyrimidine tract binding protein 1	
<b>Isotyp:</b> IgG	<b>Berechnete Masse:</b> 59 aa, 6 kDa	
<b>Immunogen Katalognummer:</b> AG3277	<b>Beobachtete Masse:</b> 59.6 kDa	

## Anwendungen

### Geprüfte Anwendungen:

IF, IHC, WB, ELISA

### In Publikationen genannte Anwendungen:

IF, IHC, IP, RIP, WB

### Getestete Reaktivität:

Human, Maus, Ratte

### Zitierte Arten:

Human, Maus, Ratte

**Hinweis-IHC: Antigendemaskierung mit TE-Puffer pH 9,0 empfohlen. (\*) Wahlweise kann die Antigendemaskierung auch mit Citratpuffer pH 6,0 erfolgen.**

### Positivkontrollen:

**WB:** A431-Zellen, BxPC-3-Zellen, HeLa-Zellen, Jurkat-Zellen, K-562-Zellen, Mauslungengewebe, Rattenherzgewebe

**IHC:** humanes Prostatakarzinomgewebe, humanes Pankreasgewebe, Maushirngewebe

**IF:** Maushirngewebe,

## Hintergrundinformationen

Polypyrimidine tract-binding protein (PTBP1), also known as heterogeneous nuclear ribonucleoprotein type I (hnRNP I), is a nuclear protein that binds pre-mRNAs in specific regions of the hnRNA-protein complexes sensitive to micrococcal nuclease. It involves in pre-mRNA splicing and the regulation of alternative splicing events. PTBP1 promotes RNA looping when bound to two separate polypyrimidine tracts in the same pre-mRNA, and promotes the binding of U2 snRNP to pre-mRNA. Besides, it can cooperate with RAVER1 to modulate switching between mutually exclusive exons during maturation of the TPM1 pre-mRNA, and represses the splicing of MAPT/Tau exon 10

## Bemerkenswerte Veröffentlichungen

Verfasser	Pubmed ID	Journal	Anwendung
Yu-Meng Sun	31387917	Blood	RIP
Juan Wang	35003358	J Cancer	WB,RIP
Li Wang	36255739	Hum Mol Genet	WB

## Lagerung

### Lagerungsbedingungen:

Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil

### Lagerungspuffer:

PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

**\*\*\* 20ul-Größen enthalten 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Ausgewählte Validierungsdaten




Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using 12582-1-AP (PTBP1 antibody), at dilution of 1:100 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




WB result of PTBP1 antibody (12582-1-AP; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PTBP1 transfected HeLa cells.



Various lysates were subjected to SDS PAGE followed by western blot with 12582-1-AP (PTBP1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 12582-1-AP (PTBP1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 12582-1-AP (PTBP1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).