

Allgemeine Informationen

Katalog-Nr.: 12813-1-AP	GenBank-Zugangsnummer: BC001135	Reinigungsmethode: Antigen-Affinitätsreinigung
Größe: 150ul , Konzentration: 400 µg/ml von Nanodrop;	GeneID (NCBI): 79054	Empfohlene Verdünnungen: IHC 1:50-1:500 IF 1:10-1:100
Wirt: Kaninchen	Vollständiger Name: transient receptor potential cation channel, subfamily M, member 8	
Isotyp: IgG	Berechnete Masse: 1104 aa, 128 kDa	
Immunogen Katalognummer: AG3483		

Anwendungen

Geprüfte Anwendungen: IF, IHC, ELISA	Positivkontrollen: IHC : humanes Leberkarzinomgewebe, humanes Kolonkarzinomgewebe, humanes Prostatakarzinomgewebe IF : SH-SY5Y-Zellen,
In Publikationen genannte Anwendungen: IF, IHC	
Getestete Reaktivität: Human	
Zitierte Arten: Human, Maus	
Hinweis-IHC: Antigenmaskierung mit TE-Puffer pH 9,0 empfohlen. (*) Wahlweise kann die Antigenmaskierung auch mit Citratpuffer pH 6,0 erfolgen.	

Hintergrundinformationen

TRPM8, also known as CMR1 (cold-menthol receptor-1), is a nonselective cation channel activated by cold and the cooling substances menthol and icilin (PMID: 11893340; 11882888). It serves as a neuronal sensor of cold temperatures. Functioning TRPM8 channels are required for behavioral responses to innocuous cool, noxious cold, injury-evoked cold hypersensitivity, and cooling-mediated analgesia (PMID: 17538622; 21438154; 16920620; 21984952). Additionally, TRPM8 has also been reported to be involved in thermoregulation and the regulation of basal tear flow (PMID: 21407809; 21076394). TRPM8 is expressed extensively in sensory nerves, human prostate and overexpressed in a variety of cancers including prostate, breast, lung, colon and skin melanomas (PMID: 11325849; 20816009).

Bemerkenswerte Veröffentlichungen

Verfasser	Pubmed ID	Journal	Anwendung
Xiyuan Zhou	36517318	J Dermatol Sci	IHC, IF
Jie Fu	36609627	Cancer Gene Ther	IHC
Xiao-Tao He	35247634	Biomaterials	IF

Lagerung


Lagerungsbedingungen:
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil
Lagerungspuffer:
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.
 Aliquotieren ist nicht notwendig bei -20°C Lagerung

***** 20ul-Größen enthalten 0.1% BSA**


For technical support and original validation data for this product please contact:
 T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
 E: proteintech@ptglab.com
 W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 12813-1-AP (TRPM8 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 12813-1-AP (TRPM8 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of SH-SY5Y cells using 12813-1-AP (TRPM8 antibody) at dilution of 1:25 and Rhodamine-Goat anti-Rabbit IgG.