

Nur für Forschungszwecke

# HDAC6 Polyklonaler Antikörper

Katalog-Nr.:12834-1-AP

Vorgestelltes Produkt

43 Publikationen



## Allgemeine Informationen

<b>Katalog-Nr.:</b> 12834-1-AP	<b>GenBank-Zugangsnummer:</b> BC013737	<b>Reinigungsmethode:</b> Antigen-Affinitätsreinigung
<b>Größe:</b> 150ul , Konzentration: 850 µg/ml von Nanodrop;	<b>GeneID (NCBI):</b> 10013	<b>Empfohlene Verdünnungen:</b> WB 1:500-1:2000 IP 0.5-4.0 ug für IP und 1:500-1:2000 für WB
<b>Wirt:</b> Kaninchen	<b>Vollständiger Name:</b> histone deacetylase 6	<b>IHC 1:20-1:200</b> <b>IF 1:50-1:500</b>
<b>Isotyp:</b> IgG	<b>Berechnete Masse:</b> 1063 aa, 114 kDa, 131 kDa	
<b>Immunogen Katalognummer:</b> AG3572	<b>Beobachtete Masse:</b> 160 kDa	

## Anwendungen

### Geprüfte Anwendungen:

IF, IHC, IP, WB, ELISA

### In Publikationen genannte Anwendungen:

CoIP, IF, IHC, IP, RIP, WB

### Getestete Reaktivität:

Human

### Zitierte Arten:

Human

**Hinweis-IHC: Antigenmaskierung mit TE-Puffer pH 9,0 empfohlen. (\*) Wahlweise kann die Antigenmaskierung auch mit Citratpuffer pH 6,0 erfolgen.**

### Positivkontrollen:

**WB :** K-562-Zellen, HeLa-Zellen, HepG2-Zellen, humanes Lebergewebe

**IP :** K-562-Zellen,

**IHC :** humanes Hodengewebe, humanes Lungenkarzinomgewebe, humanes Mammakarzinomgewebe

**IF :** HepG2-Zellen,

## Hintergrundinformationen

Histone deacetylases(HDAC) are a class of enzymes that remove the acetyl groups from the lysine residues leading to the formation of a condensed and transcriptionally silenced chromatin. At least 4 classes of HDAC were identified. HDAC6 is a member of the class II mammalian histone deacetylases. It possesses two separate putative catalytic domains. Both catalytic domains are fully functional HDACs and contribute independently to the overall activity of HDAC6 protein. A very potent NES is present at the amino-terminus of HDAC6, which was found to play an important role in regulating the shuttling of HDAC6 protein between cytoplasm and nucleus. The shuttling process may be a critical regulatory mechanism of HDAC6 function. The expression of HDAC6 is tightly linked to the state of cell differentiation. HDAC6 may participate in coordinating expression of a group of genes involved in the remodelling of chromatin during cell differentiation. HDAC6 has some splicing variants such as P114(~130kd), P131(~160kd). This antibody is a rabbit polyclonal antibody raised against residues near the C terminal of human HDAC6. The calculated molecular weight of HDAC6 is 130 kDa, but the modified the HDAC6 is about 150-160 kDa.

## Bemerkenswerte Veröffentlichungen

Verfasser	Pubmed ID	Journal	Anwendung
Shixin Gong	36117192	Cell Regen	WB
Yijing Zheng	32926854	Am J Pathol	WB
Shenli Yuan	36068616	Genome Biol	WB,IP

## Lagerung

### Lagerungsbedingungen:

Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil

### Lagerungspuffer:

PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

**\*\*\* 20ul-Größen enthalten 0.1% BSA**

For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Ausgewählte Validierungsdaten




K-562 cells were subjected to SDS PAGE followed by western blot with 12834-1-AP (HDAC6 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




IP Result of anti-HDAC6 (IP:12834-1-AP, 5ug; Detection:12834-1-AP 1:1000) with K-562 cells lysate 11000ug.




Immunohistochemical analysis of paraffin-embedded human testis using 12834-1-AP (HDAC6 antibody) at dilution of 1:50 (under 10x lens).




Immunohistochemical analysis of paraffin-embedded human testis using 12834-1-AP (HDAC6 antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of (10% Formaldehyde) fixed HepG2 cells using 12834-1-AP (HDAC6 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 12834-1-AP (HDAC6 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



WB result of HDAC6 antibody (12834-1-AP; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HDAC6 transfected HeLa cells.