

Nur für Forschungszwecke

LNx1 Polyklonaler Antikörper

Katalog-Nr.: 13124-1-AP



Allgemeine Informationen

Katalog-Nr.: 13124-1-AP	GenBank-Zugangsnummer: BC022983	Reinigungsmethode: Antigen-Affinitätsreinigung
Größe: 150ul, Konzentration: 300 µg/ml von Nanodrop und 133 µg/ml durch die Bradford-Methode mit BSA als Standard;	GeneID (NCBI): 84708	Empfohlene Verdünnungen: WB 1:500-1:2400
Wirt: Kaninchen	Vollständiger Name: Ligand of numb-protein X 1	
Isotyp: IgG	Berechnete Masse: 728 aa, 81 kDa	
Immunogen Katalognummer: AG3823	Beobachtete Masse: 70 kDa	

Anwendungen

Geprüfte Anwendungen: WB, ELISA	Positivkontrollen: WB: Maushodengewebe, Mausnierengewebe
Getestete Reaktivität: Human, Maus, Ratte	

Hintergrundinformationen

LNx1 (Ligand of Numb, protein X 1) was originally identified as a binding partner of the cell-fate determinant Numb and was reported to be involved in ubiquitylation and degradation of Numb. LNx1 contains four PDZ domains and may function as a scaffold for a multi-protein complex. LNx1 has two splicing variants, LNx1p70 and LNx1p80. LNx1p70 has the N-terminal Numb binding region and four PDZ domains, whereas LNx1p80 has an additional RING-finger domain at the N-terminus followed by a Numb-binding region and four PDZ domains. This antibody can detect both LNx1p70 and LNx1p80, around 70-80 kDa and 80-100 kDa, respectively.

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil
Lagerungspuffer:
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

***** 20ul-Größen enthalten 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



mouse testis tissue were subjected to SDS PAGE followed by western blot with 13124-1-AP (LNK1 antibody) at dilution of 1:70 incubated at room temperature for 1.5 hours.